


30 MINUTES  
A DAY ....  
GEOMETRY



# Contents



<b>INTRODUCTION</b>		vii
<b>PRETEST</b>		1
<b>LESSON 1</b>	<b>The Basic Building Blocks of Geometry</b> Explains the basic building blocks of geometry: points, lines, rays, line segments, and planes	13
<b>LESSON 2</b>	<b>Types of Angles</b> Describes right, acute, obtuse, and straight angles	23
<b>LESSON 3</b>	<b>Working with Lines</b> Describes perpendicular, transversal, parallel, and skew lines and shows how to solve problems involving them	29
<b>LESSON 4</b>	<b>Measuring Angles</b> Describes how to measure and draw a variety of angles using a protractor and how to add and subtract angle measures	37
<b>LESSON 5</b>	<b>Pairs of Angles</b> Explains the special relationships that exist between complementary, supplementary, and vertical angles	45
<b>LESSON 6</b>	<b>Types of Triangles</b> Shows how to identify scalene, isosceles, equilateral, acute, equiangular, right, and obtuse triangles	51

## CONTENTS

<b>LESSON 7</b>	<b>Congruent Triangles</b> Defines congruent triangles and shows how to prove that triangles are congruent	59
<b>LESSON 8</b>	<b>Triangles and the Pythagorean Theorem</b> Shows how the various parts of a right triangle are related and how to use the Pythagorean theorem and its converse to solve problems	67
<b>LESSON 9</b>	<b>Properties of Polygons</b> Describes plane figures such as decagons, hexagons, pentagons, and octagons and shows how to find their interior and exterior angle measurements	77
<b>LESSON 10</b>	<b>Quadrilaterals</b> Discusses the various properties of parallelograms and trapezoids and explains why they are quadrilaterals	85
<b>LESSON 11</b>	<b>Ratio, Proportion, and Similarity</b> Explains ratio, proportion, and similarity and shows how to discern if triangles are similar	93
<b>LESSON 12</b>	<b>Perimeter of Polygons</b> Explains how to find the perimeter of concave and convex polygons	103
<b>LESSON 13</b>	<b>Area of Polygons</b> Explains how to find the area of polygons	111
<b>LESSON 14</b>	<b>Surface Area of Prisms</b> Shows how to find the surface area of prisms	121
<b>LESSON 15</b>	<b>Volume of Prisms and Pyramids</b> Explains how to find the volume of prisms and pyramids	127
<b>LESSON 16</b>	<b>Working with Circles and Circular Figures</b> Explains how to find the circumference and area of circles and the surface area and volume of cylinders and spheres; includes a thorough discussion of pi	135
<b>LESSON 17</b>	<b>Coordinate Geometry</b> Introduces coordinate geometry, describes the coordinate plane, and shows how to plot points and determine distance between two points	145
<b>LESSON 18</b>	<b>The Slope of a Line</b> Defines slope and shows how to determine the slope of a line	151

## CONTENTS

<b>LESSON 19</b>	<b>The Equation of a Line</b>	157
	Describes the equation of a line and shows how to write equations of lines, including horizontal and vertical lines	
<b>LESSON 20</b>	<b>Trigonometry Basics</b>	161
	Shows how to find the trigonometric ratios sine, cosine, and tangent by using right triangles	
<b>POSTTEST</b>		169
<b>ANSWER KEY</b>		179
<b>GLOSSARY</b>		191
<b>APPENDIX A</b>	List of postulates and theorems from the book	195
<b>APPENDIX B</b>	List of additional resources readers can consult for more information on topics covered in the book	199



# Introduction



## ► **How to Use This Book**

*30 MINUTES A DAY GEOGRAPHY...* teaches basic geometry concepts in 20 self-paced lessons. The book also includes a pretest, a posttest, a glossary of mathematical terms, an appendix with postulates and theorems, and an appendix of additional resources for further study. Before you begin Lesson 1, take the pretest, which will assess your current knowledge of geometry. You'll find the answers in the answer key at the end of the book. Taking the pretest will help you determine your strengths and weaknesses in geometry. After taking the pretest, move on to Lesson 1.

Lessons 1–19 offer detailed explanations of basic geometry topics, and Lesson 20 introduces basic trigonometry. Each lesson includes example problems with step-by-step solutions. After you study the examples, you're given a chance to practice similar problems. The answers to the practice problems are in the answer key located at the back of the book. At the end of each lesson is an exercise called Skill Building until Next Time. This exercise applies the lesson's topic to an activity you may encounter in your daily life since geometry is a tool that is used to solve many real-life problems.

After you have completed all 20 lessons, take the posttest, which has the same format as the pretest, but the questions are different. Compare your scores to see how much you've improved or to identify areas in which you need more practice.

If you feel that you need more help with geometry after you complete this book, see Appendix II for additional resources to help you continue improving your geometry skills.

### ► **Make a Commitment**

Success in geometry requires effort. Make a commitment to improve your geometry skills. Work for understanding. *Why* you do a math operation is as important as *how* you do it. If you truly want to be successful, make a commitment to spend the time you need to do a good job. You can do it! When you achieve success in geometry, you will have laid a solid foundation for future challenges and success.

So sharpen your pencil and get ready to begin the pretest!

# Pretest



**B**efore you begin the first lesson, you may want to find out how much you already know and how much you need to learn. If that's the case, take the pretest in this chapter, which includes 50 multiple-choice questions covering the topics in this book. While 50 questions can't cover every geometry skill taught in this book, your performance on the pretest will give you a good indication of your strengths and weaknesses.

If you score high on the pretest, you have a good foundation and should be able to work your way through the book quickly. If you score low on the pretest, don't despair. This book will take you through the geometry concepts, step by step. If you get a low score, you may need more than 30 minutes a day to work through a lesson. However, this is a self-paced program, so you can spend as much time on a lesson as you need. *You* decide when you fully comprehend the lesson and are ready to go on to the next one.

Take as much time as you need to complete the pretest. When you are finished, check your answers in the answer key at the end of the book. Along with each answer is a number that tells you which lesson of this book teaches you about the geometry skills needed for that question.





LEARNINGEXPRESS ANSWER SHEET

- 1. (a) (b) (c) (d)
- 2. (a) (b) (c) (d)
- 3. (a) (b) (c) (d)
- 4. (a) (b) (c) (d)
- 5. (a) (b) (c) (d)
- 6. (a) (b) (c) (d)
- 7. (a) (b) (c) (d)
- 8. (a) (b) (c) (d)
- 9. (a) (b) (c) (d)
- 10. (a) (b) (c) (d)
- 11. (a) (b) (c) (d)
- 12. (a) (b) (c) (d)
- 13. (a) (b) (c) (d)
- 14. (a) (b) (c) (d)
- 15. (a) (b) (c) (d)
- 16. (a) (b) (c) (d)
- 17. (a) (b) (c) (d)

- 18. (a) (b) (c) (d)
- 19. (a) (b) (c) (d)
- 20. (a) (b) (c) (d)
- 21. (a) (b) (c) (d)
- 22. (a) (b) (c) (d)
- 23. (a) (b) (c) (d)
- 24. (a) (b) (c) (d)
- 25. (a) (b) (c) (d)
- 26. (a) (b) (c) (d)
- 27. (a) (b) (c) (d)
- 28. (a) (b) (c) (d)
- 29. (a) (b) (c) (d)
- 30. (a) (b) (c) (d)
- 31. (a) (b) (c) (d)
- 32. (a) (b) (c) (d)
- 33. (a) (b) (c) (d)
- 34. (a) (b) (c) (d)

- 35. (a) (b) (c) (d)
- 36. (a) (b) (c) (d)
- 37. (a) (b) (c) (d)
- 38. (a) (b) (c) (d)
- 39. (a) (b) (c) (d)
- 40. (a) (b) (c) (d)
- 41. (a) (b) (c) (d)
- 42. (a) (b) (c) (d)
- 43. (a) (b) (c) (d)
- 44. (a) (b) (c) (d)
- 45. (a) (b) (c) (d)
- 46. (a) (b) (c) (d)
- 47. (a) (b) (c) (d)
- 48. (a) (b) (c) (d)
- 49. (a) (b) (c) (d)
- 50. (a) (b) (c) (d)



► **Pretest**

1. Which is the correct notation for line AB?

- a.  $\overrightarrow{AB}$
- b.  $\overleftrightarrow{AB}$
- c.  $\overrightarrow{BA}$
- d.  $\overleftrightarrow{AB}$

2. Which is a correct name for this line?




- a.  $\overleftrightarrow{XZ}$
- b.  $\overleftrightarrow{ZX}$
- c.  $\overleftrightarrow{XY}$
- d.  $\overleftrightarrow{YZ}$

3. Which is not a property of a plane?


- a. is a flat surface
- b. has no thickness
- c. has boundaries
- d. has two dimensions

4. Which is a correct name for the angle?




- a.  $\angle R$
- b.  $\angle T$
- c.  $\angle TSR$
- d.  $\angle RTS$

5. Which is NOT a correct name for the angle?




- a.  $\angle RST$
- b.  $\angle TSR$
- c.  $\angle S$
- d.  $\angle R$

6. Which line is a transversal?




- a. line  $l$
- b. line  $m$
- c. line  $n$
- d. line  $r$

7. Which pairs of lines are parallel?




- a. lines  $r$  and  $s$
- b. lines  $l$  and  $m$
- c. lines  $r$  and  $n$
- d. lines  $s$  and  $l$

8. What is the best way to describe the pair of lines  $l$  and  $m$ ?




- a. perpendicular
- b. intersecting
- c. parallel
- d. skew

9. Find the measure of  $\angle AOD$ .




- a.  $55^\circ$
- b.  $90^\circ$
- c.  $75^\circ$
- d.  $130^\circ$

10. Find the measure of  $\angle BOD$ .




- a.  $55^\circ$
- b.  $90^\circ$
- c.  $75^\circ$
- d.  $130^\circ$

11. What is the measure of  $\angle RUS$ ?




- a.  $50^\circ$
- b.  $65^\circ$
- c.  $75^\circ$
- d.  $90^\circ$

12. What is the measure of  $\angle JTK$ ?




- a.  $180^\circ$
- b.  $120^\circ$
- c.  $60^\circ$
- d.  $30^\circ$

13. What is the measure of  $\angle ATJ$ ?



- a.  $30^\circ$
- b.  $60^\circ$
- c.  $120^\circ$
- d.  $180^\circ$

14. Classify the triangle by its sides.



- a. scalene
- b. isosceles
- c. equilateral
- d. none of the above

15. Classify  $\triangle ABC$  with the following measurements,  $AB = 3$ ,  $BC = 7$ , and  $AC = 7$ .

- a. scalene
- b. isosceles
- c. equilateral
- d. none of the above


16. Given  $\triangle ABC \cong \triangle XYZ$ ,  $\angle A$  corresponds to

- a.  $\angle B$ .
- b.  $\angle C$ .
- c.  $\angle X$ .
- d.  $\angle Z$ .

17. Given  $\triangle ABC \cong \triangle XYZ$ ,  $\overline{YZ}$  corresponds to


- a.  $\overline{BC}$ .
- b.  $\overline{AB}$ .
- c.  $\overline{AC}$ .
- d.  $\overline{ZY}$ .

18. Which postulate could you use to prove  $\triangle FGH \cong \triangle PQR$ ?




- a. AAA
- b. SAS
- c. ASA
- d. SSS

19. The hypotenuse of  $\triangle HLM$  is



- a.  $\overline{HL}$ .
- b.  $\overline{LM}$ .
- c.  $\overline{HM}$ .
- d.  $\overline{LH}$ .


20. A leg of  $\triangle HLM$  is




- a.  $\angle H$ .
- b.  $\angle L$ .
- c.  $\overline{HL}$ .
- d.  $\overline{HM}$ .

21. Which figure is a convex polygon?


a.




b.



c.



d.




22. Which figure is a concave polygon?


a.




b.



c.



d.



23. What is the name for a polygon with six sides?

- a. pentagon
- b. hexagon
- c. octagon
- d. decagon

24. Which of the following is not necessarily a parallelogram?

- a. quadrilateral
- b. rectangle
- c. rhombus
- d. square

25. Which of the following is NOT a quadrilateral?


- a. trapezoid
- b. parallelogram
- c. decagon
- d. square

26. Express the ratio  $\frac{JL}{JM}$  in simplest form.



- a.  $\frac{3}{7}$
- b.  $\frac{7}{3}$
- c.  $\frac{3}{10}$
- d.  $\frac{10}{3}$

27. Express the ratio  $\frac{LM}{JL}$  in simplest form.




- a.  $\frac{3}{7}$
- b.  $\frac{7}{3}$
- c.  $\frac{3}{10}$
- d.  $\frac{10}{3}$

28. Which of the following is a true proportion?

- a.  $3:7 = 5:8$
- b.  $1:2 = 4:9$
- c.  $6:3 = 10:6$
- d.  $2:5 = 4:10$

29. Find the perimeter of the polygon.



- a. 9 cm
- b. 18 cm
- c. 36 cm
- d. 72 cm

30. Find the perimeter of a square that measures 16 inches on one side.

- a. 16 in.
- b. 32 in.
- c. 64 in.
- d. not enough information

31. Find the area of a rectangle with base 7 inches and height 11 inches.

- a.  $77 \text{ in.}^2$
- b.  $36 \text{ in.}^2$
- c.  $18 \text{ in.}^2$
- d.  $154 \text{ in.}^2$


32. Find the area of a parallelogram with base 5 cm and height 20 cm.

- a.  $100 \text{ cm}^2$
- b.  $50 \text{ cm}^2$
- c.  $25 \text{ cm}^2$
- d.  $15 \text{ cm}^2$

33. Find the area of a square that measures 11 yards on an edge.

- a.  $44 \text{ yd.}^2$
- b.  $121 \text{ yd.}^2$
- c.  $22 \text{ yd.}^2$
- d.  $242 \text{ yd.}^2$


34. Which segment does not equal 3 m?




- a.  $\overline{LO}$
- b.  $\overline{HK}$
- c.  $\overline{IJ}$
- d.  $\overline{LM}$



35. Use the formula  $SA = 2(lw + wh + lh)$  to find the surface area of the prism.




- a.  $22 \text{ m}^2$   
 b.  $11 \text{ m}^2$   
 c.  $44 \text{ m}^2$   
 d.  $88 \text{ m}^2$
36. Find the volume of a prism with length 12 inches, width 5 inches, and height 8 inches.
- a.  $60 \text{ in.}^3$   
 b.  $40 \text{ in.}^3$   
 c.  $480 \text{ in.}^3$   
 d.  $96 \text{ in.}^3$
37. Find the volume of the triangular prism.



- a.  $14 \text{ cm}^3$   
 b.  $21 \text{ cm}^3$   
 c.  $42 \text{ cm}^3$   
 d.  $84 \text{ cm}^3$
38. Find the volume of a prism with base area  $50 \text{ m}^2$  and height 7 m. Use  $V = bh$ .
- a.  $17,500 \text{ m}^3$   
 b.  $57 \text{ m}^3$   
 c.  $175 \text{ m}^3$   
 d.  $350 \text{ m}^3$


39. Find the circumference of a circle with a diameter of 21 inches. Use 3.14 for  $\pi$ .
- a. 32.47 in.  
 b. 129.88 in.  
 c. 756.74 in.  
 d. 65.94 in.
40. Find the area of a circle with a diameter of 20 cm. Use 3.14 for  $\pi$ .
- a.  $628 \text{ cm}^2$   
 b.  $62.8 \text{ cm}^2$   
 c.  $314 \text{ cm}^2$   
 d.  $31.4 \text{ cm}^2$
41. Which quadrant would you graph the point  $(5, -6)$ ?
- a. I  
 b. II  
 c. III  
 d. IV

42. The coordinates for point A are



- a.  $(3, -2)$ .  
 b.  $(-2, 3)$ .  
 c.  $(-3, 2)$ .  
 d.  $(2, -3)$ .

43. The coordinates for point  $B$  are



- a.  $(3, -2)$ .  
 b.  $(-2, 3)$ .  
 c.  $(-3, 2)$ .  
 d.  $(2, -3)$ .
44. A line that points up to the right has a slope that is  
 a. positive.  
 b. negative.  
 c. zero.  
 d. undefined.
45. A vertical line has a slope that is  
 a. positive.  
 b. negative.  
 c. zero.  
 d. undefined.
46. Which of the following is a linear equation?  
 a.  $\frac{10}{x} = y$   
 b.  $3x + 2y^2 = 10$   
 c.  $3x^2 + 2y = 10$   
 d.  $3x + 2y = 10$

47. Which of the following is not a linear equation?

- a.  $x = 4$   
 b.  $y = -4$   
 c.  $\frac{1}{2}x + \frac{1}{3}y = 7$   
 d.  $\frac{2}{x} = y$

48. Which ordered pair satisfies the equation

$$3x + 4y = 12?$$

- a.  $(0, 0)$   
 b.  $(2, 2)$   
 c.  $(1, 4)$   
 d.  $(0, 3)$

49. The ratio of the opposite leg to an adjacent leg is

the trigonometric ratio

- a. sine.  
 b. cosine.  
 c. tangent.  
 d. hypotenuse.

50. The ratio of the adjacent leg to the hypotenuse is

the trigonometric ratio

- a. sine.  
 b. cosine.  
 c. tangent.  
 d. hypotenuse.



LESSON

1



# The Basic Building Blocks of Geometry

## LESSON SUMMARY

This lesson explains the basic building blocks of geometry: points, lines, rays, line segments, and planes. It also shows you the basic properties you need to understand and apply these terms.

**T**he term *geometry* is derived from the two Greek words *geo* and *metric*. It means “to measure the Earth.” The great irony is that the most basic building block in geometry, the *point*, has no measurement at all. You must accept that a point exists in order to have lines and planes because lines and planes are made up of an infinite number of points. If you do not accept the assumption that a point can exist without size or dimension, then the rest of this lesson (and book) cannot exist either. Therefore, this is a safe assumption to make! Let’s begin this lesson by looking at each of the basic building blocks of geometry.

### ► Points

A *point* has no size and no dimension; however, it indicates a definite location. Some examples are: a pencil point, a corner of a room, and the period at the end of this sentence. A series of points are what make up lines, line segments, rays, and planes. There are countless points on any one line. A point is named with an italicized uppercase letter placed next to it:

• *A*

If you want to discuss this point with someone, you would call it “point *A*.”


## ► Lines

A *line* is straight, but it has no thickness. It is an infinite set of points that extends in both directions. Imagine a straight highway with no end and no beginning; it is an example of a line. A line is named by any one italicized lowercase letter or by naming any two points on the line. If the line is named by using two points on the line, a small symbol of a line (with two arrows) is written above the two letters. For example, this line could be referred to as line  $\overleftrightarrow{AB}$  or line  $\overleftrightarrow{BA}$ :



### Practice

1. Are there more points on  $\overleftrightarrow{AB}$  than point  $A$  and point  $B$ ?
2. What are three examples of a point in everyday life?
3. Why would lines, segments, rays, or planes not exist if points do not exist?

4. Write six different names for this line. 

5. How many points are on a line?
6. Why do you think the notation for a line has two arrowheads?

► **Rays**

A *ray* is a part of a line with one endpoint. It has an infinite number of points on it. Flashlights and laser beams are good examples of rays. When you refer to a ray, you always name the endpoint first. The ray below is ray  $\overrightarrow{AB}$ , not ray  $\overrightarrow{BA}$ .



► **Line Segments**

A *line segment* is part of a line with two endpoints. It has an infinite number of points on it. A ruler and a baseboard are examples of line segments. Line segments are also named with two italicized uppercase letters, but the symbol above the letters has no arrows. This segment could be referred to as  $\overline{AB}$  or  $\overline{BA}$ :



**Practice**


7. Name two different rays with endpoint S.



8. Why is it important to name the endpoint of a ray first?

9. Why are ray  $\overrightarrow{AB}$  and ray  $\overrightarrow{BA}$  not the same?

10. Name six different line segments shown.




11. Why are arrowheads not included in line segment notation?


12. How many points are on a line segment?

## ► Planes

A *plane* is a flat surface that has no thickness or boundaries. Imagine a floor that extends in all directions as far as you can see. Here is what a plane looks like:




When you talk to someone about this plane, you could call it “plane *ABC*.” However, the more common practice is to name a plane with a single italicized capital letter (no dot after it) placed in the upper-right corner of the figure as shown here:



If you want to discuss this plane with someone, you would call it “plane *X*.”

**THE BASIC BUILDING BLOCKS OF GEOMETRY**

**THE BASIC BUILDING BLOCKS OF GEOMETRY**

FIGURE	NAME	SYMBOL	READ AS	PROPERTIES	EXAMPLES
• <i>A</i>	Point	• <i>A</i>	point <i>A</i>	<ul style="list-style-type: none"> <li>• has no size</li> <li>• has no dimension</li> <li>• indicates a definite location</li> <li>• named with an italicized uppercase letter</li> </ul>	<ul style="list-style-type: none"> <li>• pencil point</li> <li>• corner of a room</li> </ul>
$\overleftrightarrow{AB}$  $l$ $\overleftrightarrow{\quad}$	Line	$\overleftrightarrow{AB}$ or $\overleftrightarrow{BA}$	line <i>AB</i> or <i>BA</i>  or line <i>l</i>	<ul style="list-style-type: none"> <li>• is straight</li> <li>• has no thickness</li> <li>• an infinite set of points that extends in opposite directions</li> <li>• one dimension</li> </ul>	<ul style="list-style-type: none"> <li>• highway without boundaries</li> <li>• hallway without bounds</li> </ul>
$\overrightarrow{AB}$	Ray	$\overrightarrow{AB}$	ray <i>AB</i> (endpoint is always first)	<ul style="list-style-type: none"> <li>• is part of a line</li> <li>• has only one endpoint</li> <li>• an infinite set of points that extends in one direction</li> <li>• one dimension</li> </ul>	<ul style="list-style-type: none"> <li>• flashlight</li> <li>• laser beam</li> </ul>
$\overline{AB}$	Line segment	$\overline{AB}$ or $\overline{BA}$	segment <i>AB</i> or <i>BA</i>	<ul style="list-style-type: none"> <li>• is part of a line</li> <li>• has two endpoints</li> <li>• an infinite set of points</li> <li>• one dimension</li> </ul>	<ul style="list-style-type: none"> <li>• edge of a ruler</li> <li>• base board</li> </ul>
	Plane	None	plane <i>ABC</i> or plane <i>X</i>	<ul style="list-style-type: none"> <li>• is a flat surface</li> <li>• has no thickness</li> <li>• an infinite set of points that extends in all directions</li> <li>• two dimensions</li> </ul>	<ul style="list-style-type: none"> <li>• floor without boundaries</li> <li>• surface of a football field without boundaries</li> </ul>



**Practice**

- 13. A line is different from a ray because \_\_\_\_\_.
- 14. A line is similar to a ray because \_\_\_\_\_.
- 15. A ray is different from a segment because \_\_\_\_\_.
- 16. A ray is similar to a segment because \_\_\_\_\_.
- 17. A plane is different from a line because \_\_\_\_\_.

**► Working with the Basic Building Blocks of Geometry**

Points, lines, rays, line segments, and planes are very important building blocks in geometry. Without them, you cannot work many complex geometry problems. These five items are closely related to each other. You will use them in all the lessons that refer to *plane figures*—figures that are flat with one or two dimensions. Later in the book, you will study three-dimensional figures—figures that occur in space. *Space* is the set of all possible points and is three-dimensional. For example, a circle and a square are two-dimensional and can occur in a plane. Therefore, they are called *plane figures*. A sphere (ball) and a cube are examples of three-dimensional figures that occur in space, not a plane.

One way you can see how points and lines are related is to notice whether points lie in a straight line. *Collinear points* are points on the same line. *Noncollinear points* are points not on the same line. Even though you may not have heard these two terms before, you probably can correctly label the following two figures based on the sound of the names *collinear* and *noncollinear*.




Collinear points




Noncollinear points

A way to see how points and planes are related is to notice whether points lie in the same plane. For instance, see these two figures:



*Coplanar points*



*Noncoplanar points*

Again, you may have guessed which name is correct by looking at the figures and seeing that in the figure labeled *Coplanar points*, all the points are on the same plane. In the figure labeled *Noncoplanar points*, the points are on different planes.

**Practice**

- 18. Can two points be noncollinear? Why or why not?
  
- 19. Can three points be noncollinear? Why or why not?
  
- 20. Can coplanar points be noncollinear? Why or why not?
  
- 21. Can collinear points be coplanar? Why or why not?

## ► Postulates and Theorems

You need a few more tools before moving on to other lessons in this book. Understanding the terms of geometry is only part of the battle. You must also be able to understand and apply certain facts about geometry and geometric figures. These facts are divided into two categories: postulates and theorems. *Postulates* (sometimes called *axioms*) are statements accepted without proof. *Theorems* are statements that can be proved. You will be using both postulates and theorems in this book, but you will not be proving the theorems.

Geometry is the application of definitions, postulates, and theorems. Euclid is known for compiling all the geometry of his time into postulates and theorems. His masterwork, *The Elements*, written about 300 B.C., is the basis for many geometry books today. We often refer to this as Euclidean geometry.


Here are two examples of postulates:

*Postulate:* Two points determine exactly one line.

*Postulate:* Three noncollinear points determine exactly one plane.


### Practice

State whether the points are collinear.



22. A, B, C
23. A, E, F
24. B, D, F
25. A, E

State whether the points are coplanar.



**26.**  $A, B, C, E$

**27.**  $D, B, C, E$

**28.**  $B, C, E, F$

**29.**  $A, B, E$

State whether the following statements are true or false.

**30.**  $\overleftrightarrow{XY}$  and  $\overleftrightarrow{YX}$  are the same line.

**31.**  $\overrightarrow{XY}$  and  $\overrightarrow{YX}$  are the same ray.

**32.**  $\overline{XY}$  and  $\overline{YX}$  are the same segment.

**33.** Any four points  $W, X, Y,$  and  $Z$  must lie in exactly one plane.

Draw and label a figure to fit each description, if possible. Otherwise, state “not possible.”

**34.** four collinear points

**35.** two noncollinear points

**36.** three noncoplanar points

### Skill Building until Next Time

When you place a four-legged table on a surface and it wobbles, then one of the legs is shorter than the other three. You can use three legs to support something that must be kept steady. Why do you think this is true?

Throughout the day, be on the lookout for some three-legged objects that support this principle. Examples include a camera tripod and an artist's easel. The bottoms of the three legs must represent three noncollinear points and determine exactly one plane.

LESSON

2




# Types of Angles

## LESSON SUMMARY

This lesson will teach you how to classify and name several types of angles. You will also learn about opposite rays.

**P**eople often use the term *angle* in everyday conversations. For example, they talk about camera angles, angles for pool shots and golf shots, and angles for furniture placement. In geometry, an *angle* is formed by two rays with a common endpoint. The symbol used to indicate an angle is  $\angle$ . The two rays are the *sides* of the angle. The common endpoint is the *vertex* of the angle. In the following figure, the sides are  $\overrightarrow{RD}$  and  $\overrightarrow{RY}$ , and the vertex is  $R$ .



### ► Naming Angles


People call you different names at different times. Sometimes, you are referred to by your full name, and other times, only by your first name or maybe even a nickname. These different names don't change who you are—just the way in which others refer to you. You can be named differently according to the setting you're in. For example,

## TYPES OF ANGLES

you may be called your given name at work, but your friends might call you by a nickname. There can be confusion sometimes when these names are different.


Similar to you, an angle can be named in different ways. The different ways an angle can be named may be confusing if you do not understand the logic behind the different methods of naming.

If three letters are used to name an angle, then the middle letter always names the vertex. If the angle does not share the vertex point with another angle, then you can name the angle with only the letter of the vertex. If a number is written inside the angle that is not the angle measurement, then you can name the angle by that number. You can name the following angle any one of these names:  $\angle WET$ ,  $\angle TEW$ ,  $\angle E$ , or  $\angle 1$ .




### Practice


1. Name the vertex and sides of the angle.



2. Name the vertex and sides of the angle.




3. Name the angle in four different ways.




**Right Angles**

Angles that make a square corner are called *right angles*. In drawings, the following symbol is used to indicate a right angle:




**Straight Angles**

Opposite rays are two rays with the same endpoint and that form a line. They form a *straight angle*. In the following figure,  $\vec{HD}$  and  $\vec{HS}$  are opposite rays.



**Practice**


Use the following figure to answer practice problems 4–7.



4. Name two right angles.
5. Name a straight angle.
6. Name a pair of opposite rays.
7. Why is  $\angle O$  an incorrect name for any of the angles shown?



Use the following figure to answer practice problems 8 and 9.



8. Are  $\overrightarrow{LM}$  and  $\overrightarrow{MN}$  opposite rays? Why or why not?



9. If two rays have the same endpoints, do they have to be opposite rays? Why or why not?

### ► Classifying Angles

Angles are often classified by their measures. The degree is the most commonly used unit for measuring angles. One full turn, or a circle, equals  $360^\circ$ .



#### Acute Angles

An *acute angle* has a measure between  $0^\circ$  and  $90^\circ$ . Here are two examples of acute angles:





#### Right Angles

A *right angle* has a  $90^\circ$  measure. The corner of a piece of paper will fit exactly into a right angle. Here are two examples of right angles:




#### Obtuse Angles

An *obtuse angle* has a measure between  $90^\circ$  and  $180^\circ$ . Here are two examples of obtuse angles:




**Straight Angles**

A straight angle has a  $180^\circ$  measure. This is an example of a straight angle ( $\angle ABC$  is a straight angle):



**Practice**

Use the following figure to answer practice problems 10–13.



- 10. Name three acute angles.
- 11. Name three obtuse angles.
- 12. Name two straight angles.
- 13. If  $\angle MON$  measures  $27^\circ$ , then  $\angle JOK$  measures \_\_\_\_\_ degrees.

Complete each statement.

- 14. An angle with measure  $90^\circ$  is called a(n) \_\_\_\_\_ angle.
- 15. An angle with measure  $180^\circ$  is called a(n) \_\_\_\_\_ angle.
- 16. An angle with a measure between  $0^\circ$  and  $90^\circ$  is called a(n) \_\_\_\_\_ angle.
- 17. An angle with a measure between  $90^\circ$  and  $180^\circ$  is called a(n) \_\_\_\_\_ angle.

## TYPES OF ANGLES

Questions 18–21 list the measurement of an angle. Classify each angle as acute, right, obtuse, or straight.

18.  $10^\circ$


20.  $98^\circ$

19.  $180^\circ$


21.  $90^\circ$

You may want to use a corner of a piece of scratch paper for questions 22–25. Classify each angle as acute, right, obtuse, or straight.


22.




23.



24.



25.



### Skill Building until Next Time

In conversational English, the word *acute* can mean sharp and the word *obtuse* can mean dull or not sharp. How can you relate these words with the drawings of acute and obtuse angles?

L E S S O N

# 3



## Working with Lines

### LESSON SUMMARY

This lesson introduces you to perpendicular, transversal, parallel, and skew lines. The angles formed by a pair of parallel lines and a transversal are also explained.


**B**oth intersecting and nonintersecting lines surround you. Most of the time you do not pay much attention to them. In this lesson, you will focus on two different types of intersecting lines: transversals and perpendicular lines. You will also study nonintersecting lines: parallel and skew lines. You will learn properties about lines that have many applications to this lesson and throughout this book. You'll soon start to look at the lines around you with a different point of view.

### ► Intersecting Lines


On a piece of scratch paper, draw two straight lines that cross. Can you make these straight lines cross at more than one point? No, you can't, because intersecting lines cross at only one point (unless they are the same line). The point where they cross is called a *point of intersection*. They actually share this point because it is on both lines. Two special types of intersecting lines are called *transversals* and *perpendicular lines*.

### Transversals

A *transversal* is a line that intersects two or more other lines, each at a different point. In the following figure, line  $t$  is a transversal, line  $s$  is not.




The prefix *trans* means to cross. In the previous figure, you can see that line  $t$  cuts across the two lines  $m$  and  $n$ . Line  $m$  is a transversal for lines  $s$  and  $t$ . Also, line  $n$  is a transversal across lines  $s$  and  $t$ . Line  $s$  crosses lines  $m$  and  $n$  at the same point (their point of intersection); therefore, line  $s$  is not a transversal. A transversal can cut across parallel as well as intersecting lines, as shown here:



### Practice


Use the following figure to answer questions 1–4.




1. Is line  $d$  a transversal? Why or why not?
2. Is line  $y$  a transversal? Why or why not?
3. Is line  $t$  a transversal? Why or why not?
4. Is line  $r$  a transversal? Why or why not?

### Perpendicular Lines

*Perpendicular lines* are another type of intersecting lines. Everyday examples of perpendicular lines include the horizontal and vertical lines of a plaid fabric and the lines formed by panes in a window. Perpendicular lines meet to form right angles. Right angles always measure  $90^\circ$ . In the following figure, lines  $x$  and  $y$  are perpendicular:



The symbol “ $\perp$ ” means perpendicular. You could write  $x \perp y$  to show these lines are perpendicular. Also, the symbol that makes a square in the corner where lines  $x$  and  $y$  meet indicates a right angle. In geometry, you shouldn’t assume anything without being told. Never assume a pair of lines are perpendicular without one of these symbols. A transversal *can* be perpendicular to a pair of lines, but it does not *have* to be. In the following figure, line  $t$  is perpendicular to both line  $l$  and line  $m$ .




### Practice

State whether the following statements are true or false.


- \_\_\_\_\_ **5.** Perpendicular lines always form right angles.
- \_\_\_\_\_ **6.** The symbol “ $\perp$ ” means parallel.
- \_\_\_\_\_ **7.** Transversals must always be parallel.

### Nonintersecting Lines

If lines do not intersect, then they are either parallel or skew.




Lines  $l$  and  $m$  are parallel.  
 Lines  $l$  and  $m$  are coplanar lines.  
 Lines  $l$  and  $m$  do not intersect.



Lines  $j$  and  $k$  are skew.  
 Lines  $j$  and  $k$  are noncoplanar lines.  
 Lines  $j$  and  $k$  do not intersect.

The symbol “ $\parallel$ ” means parallel. So you can abbreviate the sentence, “Lines  $l$  and  $m$  are parallel,” by writing “ $l \parallel m$ .” Do not assume a pair of lines are parallel unless it is indicated. Arrowheads on the lines in a figure indicate that the lines are parallel. Sometimes, double arrowheads are necessary to differentiate two sets of parallel lines, as shown in the following figure:



Everyday examples of parallel lines include rows of crops on a farm and railroad tracks. An example of skew lines is the vapor trails of a northbound jet and a westbound jet flying at different altitudes. One jet would pass over the other, but their paths would not cross.

### Practice

Complete the sentences with the correct word: *always*, *sometimes*, or *never*.


8. Parallel lines are \_\_\_\_\_ coplanar.
9. Parallel lines \_\_\_\_\_ intersect.
10. Parallel lines are \_\_\_\_\_ cut by a transversal.

11. Skew lines are \_\_\_\_\_ coplanar.

12. Skew lines \_\_\_\_\_ intersect.


**Angles Formed by Parallel Lines and a Transversal**

If a pair of parallel lines are cut by a transversal, then eight angles are formed. In the following figure, line  $l$  is parallel to line  $m$  and line  $t$  is a transversal forming angles 1–8. Angles 3, 4, 5, and 6 are inside the parallel lines and are called *interior* angles. Angles 1, 2, 7, and 8 are outside the parallel lines and are called *exterior* angles.




**Alternate Interior Angles**

Alternate interior angles are interior angles on opposite sides of the transversal. In the previous figure, angles 3 and 5 and angles 4 and 6 are examples of alternate interior angles. To spot alternate interior angles, look for a *Z-shaped* figure, as shown in the following figures:



**Same-Side Interior Angles**


Same-side interior angles are interior angles on the same side of the transversal. To spot same-side interior angles, look for a *U-shaped* figure in various positions, as shown in the following examples:






### Corresponding Angles

Corresponding angles are so named because they appear to be in corresponding positions in relation to the two parallel lines. Examples of corresponding angles in the following figure are angles 1 and 5, 4 and 8, 2 and 6, and 3 and 7.




To spot corresponding angles, look for an *F-shaped* figure in various positions, as shown in these examples:



### Angles Formed by Nonparallel Lines and a Transversal

Even when lines cut by a transversal are not parallel, you still use the same terms to describe angles, such as corresponding, alternate interior, and same-side interior angles. For example, look at the following figure:



- $\angle 1$  and  $\angle 5$  are corresponding angles.
- $\angle 3$  and  $\angle 5$  are alternate interior angles.
- $\angle 4$  and  $\angle 5$  are same-side interior angles.

### ► Postulates and Theorems


As mentioned in Lesson 1, *postulates* are statements of fact that we accept without proof. *Theorems* are statements of fact that can be proved. You will apply both types of facts to problems in this book. Some geometry books teach formal proofs of theorems. Although you will not go through that process in this book, you will still use both postulates and theorems for applications. There are some important facts you need to know about the special pairs of angles formed by two parallel lines and a transversal. If you noticed that some of those pairs of angles appear to have equal measure, then you are on the right track. The term used for two angles with equal measure is *congruent*.

Another important fact you should know is that a pair of angles whose sum is  $180^\circ$  is called *supplementary*. Here are the theorems and the postulate that you will apply to a figure in the next few practice exercises.

*Postulate:* If two parallel lines are cut by a transversal, then corresponding angles are congruent.  
*Theorem:* If two parallel lines are cut by a transversal, then alternate interior angles are congruent.  
*Theorem:* If two parallel lines are cut by a transversal, then same-side interior angles are supplementary.

### Practice

In the following figure,  $m \parallel n$  and  $s \parallel t$ . For questions 13–16, (a) state the special name for each pair of angles (alternate interior angles, corresponding angles, and same-side interior angles), then (b) tell if the angles are congruent or supplementary.



**13.**  $\angle 2$  and  $\angle 10$   
 a. \_\_\_\_\_ b. \_\_\_\_\_

**15.**  $\angle 13$  and  $\angle 15$   
 a. \_\_\_\_\_ b. \_\_\_\_\_

**14.**  $\angle 6$  and  $\angle 7$   
 a. \_\_\_\_\_ b. \_\_\_\_\_

**16.**  $\angle 11$  and  $\angle 14$   
 a. \_\_\_\_\_ b. \_\_\_\_\_

For 17–20, use the measure of the given angle to find the missing angle.

17.  $m\angle 2 = 100^\circ$ , so  $m\angle 7 = \_\?_\_$

18.  $m\angle 8 = 71^\circ$ , so  $m\angle 12 = \_\?_\_$

19.  $m\angle 5 = 110^\circ$ , so  $m\angle 7 = \_\?_\_$

20.  $m\angle 2 = 125^\circ$ , so  $m\angle 11 = \_\?_\_$

Complete the statement for practice problems 21–26.

21. Alternate interior angles are similar to corresponding angles because \_\_\_\_\_.

22. Alternate interior angles differ from corresponding angles because \_\_\_\_\_.

23. Same-side interior angles are similar to alternate interior angles because \_\_\_\_\_.

24. Same-side interior angles differ from alternate interior angles because \_\_\_\_\_.

25. Same-side interior angles are similar to corresponding angles because \_\_\_\_\_.

26. Same-side interior angles differ from corresponding angles because \_\_\_\_\_.

### Skill Building until Next Time

Look around your neighborhood for a building under construction. Are the floor beams the same distance apart at the front of the building as they are at the back? Why is this important? If they were not the same distance apart, how would this affect the floor? Of course, the floor would sag. Building codes specify how far apart floor beams can be based on safety codes. You can see the importance of parallel lines in construction.

LESSON

4




# Measuring Angles

## LESSON SUMMARY

This lesson focuses on using the protractor to measure and draw angles. You will also add and subtract angle measures.

**A**n instrument called a *protractor* can be used to find the measure of degrees of an angle. Most protractors have two scales, one reading from left to right, and the other, from right to left. Estimating whether an angle is acute, right, obtuse, or straight before measuring will help you choose the correct scale. Here is an example of a protractor:




Carefully line up your protractor by placing the center point of the protractor scale on the vertex of the angle, which is the place where both sides of the angle meet. If the sides of the angle do not reach the scale, extend them. Choose the scale that has zero at one side of the angle. Read the measure of the angle. Check to see if your measurement and estimate agree. When measuring an angle, it is not necessary to have one of the rays passing

through zero on the protractor scale. The angle could be measured by counting. Putting the ray on zero simply makes the counting easier.

**Practice**

Using the following figure, find the measure of each angle.



\_\_\_\_\_ 1.  $\angle AOB$

\_\_\_\_\_ 2.  $\angle AOC$

\_\_\_\_\_ 3.  $\angle AOD$

\_\_\_\_\_ 4.  $\angle AOE$

\_\_\_\_\_ 5.  $\angle BOC$

\_\_\_\_\_ 6.  $\angle BOD$

\_\_\_\_\_ 7.  $\angle BOE$


\_\_\_\_\_ 8.  $\angle BOF$

\_\_\_\_\_ 9.  $\angle COD$


\_\_\_\_\_ 10.  $\angle COE$

**► Drawing Angles**

You can use a protractor to draw an angle of a given size. First, draw a ray and place the center point of the protractor on the endpoint of the ray. Align the ray with the base line of the protractor. Locate the degree of the angle you wish to draw. Make a dot at that point and connect it to the endpoint of the ray.



The resulting angle will have the correct degree of measurement:



**Practice**

Use a protractor to draw angles with the given measures.

**11.**  $45^\circ$

**14.**  $125^\circ$



**12.**  $75^\circ$

**15.**  $32^\circ$


**13.**  $100^\circ$

### ► Adding and Subtracting Angle Measures

The following figures suggest that you can add and subtract angle measures:



*Adjacent* angles are two angles in the same plane that have the same vertex and a common side but do not have any interior points in common.




In the previous figure,  $\angle 1$  and  $\angle 2$  are adjacent angles. Angles  $1$  and  $\angle XYZ$  are not adjacent.

Use a protractor to measure  $\angle 1$ ,  $\angle 2$ , and  $\angle XYZ$ . What relationship do you notice among the measures of the three angles? You should find that the measure of  $\angle 1$  plus the measure of  $\angle 2$  equals the measure of  $\angle XYZ$ . The letter  $m$  is used before the angle symbol to represent the word *measure*. For example,  $m\angle 1 + m\angle 2 = m\angle XYZ$ . If you draw another pair of adjacent angles and measure the angles, will the relationship be the same? Try it and see.


### Angle Addition Postulate

(1) If point  $B$  is in the interior of  $\angle AOC$ , then



$$m\angle AOB + m\angle BOC = m\angle AOC.$$

(2) If  $\angle AOC$  is a straight angle, then




$$m\angle AOB + m\angle BOC = 180^\circ.$$

**Examples:** Find the measure of each angle.

(a)  $m\angle 1 = 40^\circ$

$m\angle 2 = 75^\circ$

$m\angle RSW = \underline{\hspace{2cm}}$




$$m\angle 1 + m\angle 2 = m\angle RSW$$

$$m\angle RSW = 115^\circ$$

(b)  $m\angle DEG = 35^\circ$

$m\angle GEF = 145^\circ$

$m\angle DEF = \underline{\hspace{2cm}}$



$$m\angle DEG + m\angle GEF = m\angle DEF$$

$$m\angle DEF = 180^\circ$$




**Practice**

Find the measure of each angle.

**16.**  $m\angle KJB = 60^\circ$

$m\angle BJT = 40^\circ$


$m\angle KJT = \underline{\hspace{2cm}}$



**17.**  $m\angle DOS = 120^\circ$


$m\angle DOH = 70^\circ$

$m\angle HOS = \underline{\hspace{2cm}}$




**18.**  $m\angle RPM = 42^\circ$


$m\angle MPS = \underline{\hspace{2cm}}$



19.  $m\angle GEO = 100^\circ$   
 $m\angle GEM = 60^\circ$   
 $m\angle MEO = \underline{\hspace{2cm}}$




Use the following figure to answer problems 20–25.



20.  $m\angle BPC = \underline{\hspace{2cm}}$   
 21.  $m\angle CPD = \underline{\hspace{2cm}}$   
 22.  $m\angle EPB = \underline{\hspace{2cm}}$   
 23.  $m\angle CPE = \underline{\hspace{2cm}}$   
 24.  $m\angle APE = \underline{\hspace{2cm}}$   
 25.  $m\angle DPE = \underline{\hspace{2cm}}$

## MEASURING ANGLES

Use the following figure for practice problems 26–29.



The U.S. Army uses a unit of angle measure called a *mil*. A mil is defined as  $\frac{1}{6,400}$  of a circle. The protractor is marked in mils. Find the measure in mils of each of the following angles.

- \_\_\_\_\_ **26.**  $90^\circ$  angle
- \_\_\_\_\_ **27.**  $135^\circ$  angle
- \_\_\_\_\_ **28.**  $45^\circ$  angle
- \_\_\_\_\_ **29.**  $180^\circ$  angle

### Skill Building until Next Time

Look around your environment for angles and estimate their measurements. For example, estimate the size of an angle created by leaving your door open part-way. You may also want additional practice using your protractor. If so, practice drawing several angles of different measurements.

LESSON

# 5



## Pairs of Angles

### LESSON SUMMARY

In this lesson, you will learn how to use the relationships among three special pairs of angles: complementary angles, supplementary angles, and vertical angles.

**W**hen two angles come together to form a  $90^\circ$  angle, the angle can be very useful for solving a variety of problems. Angles measuring  $90^\circ$  are essential in construction and sewing, just to name two areas.


Pairs of angles whose measures add up to  $90^\circ$  are called *complementary* angles. If you know the measurement of one angle, then you can find the measurement of the other.

When two angles come together to form a  $180^\circ$  angle, or a straight line, the applications are endless. These pairs of angles are called *supplementary* angles. You can find the measurement of one angle if the measurement of the other is given.

When two lines intersect, two pairs of *vertical* angles are always formed. You can see pairs of intersecting lines in fabric patterns, construction sites, and on road maps. If you know the measurement of one of these angles, you can find the measurement of the other three angles.

### ► Complementary Angles

Two angles are complementary angles if and only if the sum of their measure is  $90^\circ$ . Each angle is a complement of the other. To be complementary, a pair of angles do not need to be adjacent.



In the previous figure,  $\angle AOB$  and  $\angle BOC$  are a pair of complementary angles.  $\angle AOB$  and  $\angle D$  are also a pair of complementary angles.  $\angle AOB$  is a complement of  $\angle BOC$  and  $\angle D$ .

### Practice

Find the measure of a complement of  $\angle 1$  for each of the following measures of  $\angle 1$ .

\_\_\_\_\_ 1.  $m\angle 1 = 44^\circ$

\_\_\_\_\_ 2.  $m\angle 1 = 68^\circ$


\_\_\_\_\_ 3.  $m\angle 1 = 80^\circ$

\_\_\_\_\_ 4.  $m\angle 1 = 25^\circ$

\_\_\_\_\_ 5.  $m\angle 1 = 3^\circ$

### ► Supplementary Angles

Two angles are supplementary angles if and only if the sum of their measures is  $180^\circ$ . Each angle is a supplement of the other. Similar to complementary angles, supplementary angles do not need to be adjacent.



In the previous figure,  $\angle XOY$  and  $\angle YOZ$  are supplementary angles.  $\angle XOY$  and  $\angle W$  are also supplementary angles.  $\angle XOY$  is a supplement of  $\angle YOZ$  and  $\angle W$ .

**Practice**

Find the measure of a supplement of  $\angle 2$  for each of the following measures of  $\angle 2$ .

\_\_\_\_\_ **6.**  $m\angle 2 = 78^\circ$

\_\_\_\_\_ **7.**  $m\angle 2 = 130^\circ$


\_\_\_\_\_ **8.**  $m\angle 2 = 60^\circ$

\_\_\_\_\_ **9.**  $m\angle 2 = 155^\circ$

\_\_\_\_\_ **10.**  $m\angle 2 = 1^\circ$

**► Vertical Angles**

Vertical angles are two angles whose sides form two pairs of opposite rays. When two lines intersect, they form two pairs of vertical angles.




In the previous figure, the two pairs of vertical angles are  $\angle 1$  and  $\angle 3$ , as well as  $\angle 2$  and  $\angle 4$ . Also,  $\angle 1$  and  $\angle 2$  are supplementary angles. Since  $\angle 1$  and  $\angle 3$  are both supplements to the same angle, they are congruent; in other words, they have the same measurement.

*Vertical Angles theorem: Vertical angles are congruent.*

**Practice**

Use the following figure to answer practice problems 11–13.



- 11. Name two supplement angles of  $\angle DOE$ .
- 12. Name a pair of complementary angles.
- 13. Name two pairs of vertical angles.


State whether the following statements are true or false.

- \_\_\_\_\_ 14. Complementary angles must be acute.
- \_\_\_\_\_ 15. Supplementary angles must be obtuse.
- \_\_\_\_\_ 16. Two acute angles can be supplementary.
- \_\_\_\_\_ 17. A pair of vertical angles can be complementary.
- \_\_\_\_\_ 18. A pair of vertical angles can be supplementary.
- \_\_\_\_\_ 19. Vertical angles must have the same measure.
- \_\_\_\_\_ 20. Complementary angles can be adjacent.
- \_\_\_\_\_ 21. Supplementary angles can be adjacent.
- \_\_\_\_\_ 22. Any two right angles are supplementary.
- \_\_\_\_\_ 23. Two acute angles are always complementary.

\_\_\_\_\_ **24.** An acute and an obtuse angle are always supplementary.

\_\_\_\_\_ **25.** Congruent complementary angles each measure  $50^\circ$ .

Use the following figure to answer practice problem 26.



**26.** A common error is assuming that any pair of angles that are “across from each other” are vertical. In this figure,  $\angle 1$  and  $\angle 3$  are vertical angles. Angles 2 and 4 are not vertical angles. Name three other pairs of non-adjacent angles that are also not vertical.

### Skill Building until Next Time

Find a pair of intersecting lines. You may use the lines in a floor tile, a fabric design, a piece of furniture, or you could draw your own. Measure the four angles that the intersecting line forms to confirm the Vertical Angles theorem.





LESSON

6



# Types of Triangles

## LESSON SUMMARY


In this lesson, you will learn how to classify triangles according to the length of their sides and their angle measurements.

It would be difficult to name an occupation where classifying triangles is a required skill; however, it is a skill that will help you solve complex geometry problems. Each of the triangles discussed in this lesson has special properties that will help you solve problems.


### ► Classification by Sides

You can classify triangles by the lengths of their sides. On the next page are three examples of special triangles called *equilateral*, *isosceles*, and *scalene*.


## TYPES OF TRIANGLES



Isosceles  
two  $\cong$  sides



Equilateral  
three  $\cong$  sides




Scalene  
no sides congruent

To show that two or more sides of a triangle have the same measurement, a hatch mark is made through the congruent sides. Sometimes, two hatch marks are made on each congruent side, and sometimes, three hatch marks are made on each congruent side. You can match up the number of hatch marks to find which sides are congruent. You'll see these hatch marks in most geometry books. The symbol for congruent is  $\cong$ .


### Practice

Classify each triangle shown or described as equilateral, isosceles, or scalene.


1.




2.



3.



4.



5.  $\triangle ABC$  with  $AB = 10$ ,  $BC = 10$ , and  $AC = 8$ .


6.  $\triangle DEF$  with  $DE = 6$ ,  $EF = 8$ , and  $DF = 10$ .

7.  $\triangle XYZ$  with  $XY = 7$ ,  $YZ = 7$ , and  $XZ = 7$ .

### Isosceles Triangles

Isosceles triangles are important geometric figures to understand. Some geometry books define isosceles as having *at least* two congruent sides. For our purposes, we will define isosceles as having exactly two congruent sides. Did you know that the parts of an isosceles triangle have special names? The two congruent sides of an isosceles triangle are called the *legs*. The angle formed by the two congruent sides is called the *vertex angle*. The other two angles are called the *base angles*. And finally, the side opposite the vertex angle is called the *base*.

**Example:**




legs:  $\overline{AC}$  and  $\overline{AB}$   
vertex angle:  $\angle A$

base angles:  $\angle B$  and  $\angle C$   
base:  $\overline{BC}$

### Practice


Name the legs, vertex angle, base angles, and base of the isosceles triangles.

8.




legs: \_\_\_\_\_  
vertex angle: \_\_\_\_\_  
base angles: \_\_\_\_\_  
base: \_\_\_\_\_

9.



legs: \_\_\_\_\_  
vertex angle: \_\_\_\_\_  
base angles: \_\_\_\_\_  
base: \_\_\_\_\_


10.




legs: \_\_\_\_\_  
 vertex angle: \_\_\_\_\_  
 base angles: \_\_\_\_\_  
 base: \_\_\_\_\_

► **Classification by Angles**


You can also classify triangles by the measurements of their angles. Here are four examples of special triangles. They are called *acute*, *equiangular*, *right*, and *obtuse*.




Acute  
three acute angles



Equiangular  
three  $\cong$  angles



Right  
one right angle




Obtuse  
one obtuse angle

To show that two or more angles of a triangle have the same measurement, a small curve is made in the congruent angles. You can also use two small curves to show that angles are congruent.


**Practice**

Classify each triangle shown or described as acute, right, obtuse, or equiangular.


**11.**




**12.**



**13.**



**14.**



**15.**  $\triangle MNO$  with  $m\angle M = 130^\circ$ ,  $m\angle N = 30^\circ$ , and  $m\angle O = 20^\circ$ .

**16.**  $\triangle RST$  with  $m\angle R = 80^\circ$ ,  $m\angle S = 45^\circ$ , and  $m\angle T = 55^\circ$ .

**17.**  $\triangle GHI$  with  $m\angle G = 20^\circ$ ,  $m\angle H = 70^\circ$ , and  $m\angle I = 90^\circ$ .

State whether the following statements are true or false.

\_\_\_\_\_ **18.** An equilateral triangle is also an equiangular triangle.

\_\_\_\_\_ **19.** An isosceles triangle may also be a right triangle.

\_\_\_\_\_ **20.** It is possible for a scalene triangle to be equiangular.

\_\_\_\_\_ **21.** It is possible for a right triangle to be equilateral.

\_\_\_\_\_ **22.** It is possible for an obtuse triangle to be isosceles.

\_\_\_\_\_ **23.** An equilateral triangle is also an acute triangle.

\_\_\_\_\_ **24.** A triangle can have two obtuse angles.


\_\_\_\_\_ **25.** A triangle cannot have more than one right angle.




**Skill Building until Next Time**

A triangle is called a rigid figure because the length of at least one side of a triangle must be changed before its shape changes. This is an important quality in construction. Notice the triangles you can see in floor and roof trusses.


To prove the strength of a triangle, place two glasses about four inches apart. Bridge the cups with a dollar bill.




Then try to balance a box of toothpicks on the dollar bill without the box of toothpicks touching the glasses.



Fold the dollar bill like a fan.



Now try to balance the box of toothpicks on the folded dollar bill.



When you look at the folded dollar bill from the side, what shape do you see?



The view of the dollar bill from the side shows several triangles.

LESSON




# Congruent Triangles

## LESSON SUMMARY

This lesson will help you identify corresponding parts of congruent triangles and to name the postulate or theorem that shows that two triangles are congruent.

**C**ongruent triangles are commonly used in the construction of quilts, buildings, and bridges. Congruent triangles are also used to estimate inaccessible distances; for example, the width of a river or the distance across a canyon. In this lesson, you will learn simple ways to determine whether two triangles are congruent.

When you buy floor tiles, you get tiles that are all the same shape and size. One tile will fit right on top of another. In geometry, you would say one tile is *congruent* to another tile. Similarly, in the following figure,  $\triangle ABC$  and  $\triangle XYZ$  are congruent. They have the same size and shape.



## CONGRUENT TRIANGLES

Imagine sliding one triangle over to fit on top of the other triangle. You would put point  $A$  on point  $X$ ; point  $B$  on point  $Y$ ; and point  $C$  on point  $Z$ . When the vertices are matched in this way,  $\angle A$  and  $\angle X$  are called *corresponding angles*, and  $\overline{AB}$  and  $\overline{XY}$  are called *corresponding sides*.


Corresponding angles and corresponding sides are often referred to as corresponding parts of the triangles. In other words, you could say Corresponding Parts of Congruent Triangles are Congruent (CPCTC). This statement is often referred to by the initials CPCTC.

When  $\triangle ABC$  is congruent to  $\triangle XYZ$ , you write  $\triangle ABC \cong \triangle XYZ$ . This means that all of the following are true:

$$\begin{array}{lll} \angle A \cong \angle X & \angle B \cong \angle Y & \angle C \cong \angle Z \\ \overline{AB} \cong \overline{XY} & \overline{BC} \cong \overline{YZ} & \overline{AC} \cong \overline{XZ} \end{array}$$

Suppose instead of writing  $\triangle ABC \cong \triangle XYZ$ , you started to write  $\triangle CAB \cong$  \_\_\_\_\_. Since you started with  $C$  to name the first triangle, you must start with the corresponding letter,  $Z$ , to name the second triangle. Corresponding parts are named in the same order. If you name the first triangle  $\triangle CAB$ , then the second triangle must be named  $\triangle ZXY$ . In other words,  $\triangle CAB \cong \triangle ZXY$ .

**Example:** Name the corresponding angles and corresponding sides.




**Solution:**

Corresponding angles:  $\angle R$  and  $\angle E$ ;  $\angle S$  and  $\angle F$ ;  $\angle T$  and  $\angle G$

Corresponding sides:  $\overline{RS}$  and  $\overline{EF}$ ;  $\overline{ST}$  and  $\overline{FG}$ ;  $\overline{RT}$  and  $\overline{EG}$

### Practice

For practice problems 1–6, complete each statement; given  $\triangle JKM \cong \triangle PQR$ .



1.  $\angle M$  corresponds to  $\angle$ \_\_\_\_\_.

4.  $\overline{JK}$  corresponds to \_\_\_\_\_.


2.  $\angle P$  corresponds to  $\angle$ \_\_\_\_\_.

5.  $\overline{RQ}$  corresponds to \_\_\_\_\_.

3.  $\angle Q$  corresponds to  $\angle$ \_\_\_\_\_.

6.  $\overline{PR}$  corresponds to \_\_\_\_\_.

For practice problems 7–10, complete each statement, given  $\triangle GFH \cong \triangle JFH$ .



7.  $\triangle GFH \cong \triangle$ \_\_\_\_\_

9.  $\triangle FHJ \cong \triangle$ \_\_\_\_\_

8.  $\triangle HJF \cong \triangle$ \_\_\_\_\_


10.  $\triangle JFH \cong \triangle$ \_\_\_\_\_

### ► Side-Side-Side (SSS) Postulate

If you have three sticks that make a triangle and a friend has identical sticks, would it be possible for each of you to make different-looking triangles? No, it is impossible to do this. A postulate of geometry states this same idea. It is called the *side-side-side postulate*.

*Side-Side-Side postulate:* If three sides of one triangle are congruent to three sides of another triangle, then the two triangles are congruent.

Take a look at the following triangles to see this postulate in action:




The hatch marks on the triangles show which sides are congruent to which in the two triangles. For example,  $\overline{AC}$  and  $\overline{RT}$  both have one hatch mark, which shows that these two segments are congruent.  $\overline{BC}$  is congruent to  $\overline{ST}$ , as shown by the two hatch marks, and  $\overline{AB}$  and  $\overline{RS}$  are congruent as shown by the three hatch marks.

Since the markings indicate that the three pairs of sides are congruent, you can conclude that the three pairs of angles are also congruent. From the definition of congruent triangles, it follows that all six parts of  $\triangle ABC$  are congruent to the corresponding parts of  $\triangle RST$ .

**Practice**

Use the following figure to answer questions 11–15.



11.  $\overline{RS}$  corresponds to \_\_\_\_\_.

12.  $\overline{TS}$  corresponds to \_\_\_\_\_.

13.  $\overline{RV}$  corresponds to \_\_\_\_\_.

14. Is  $\triangle RTS \cong \triangle RVS$ ?


15. Is  $\triangle RSV \cong \triangle RTS$ ?

**► Side-Angle-Side (SAS) Postulate**

If you put two sticks together at a certain angle, there is only one way to finish forming a triangle. Would it be possible for a friend to form a different-looking triangle if he or she started with the same two lengths and the same angle? No, it would be impossible. Another postulate of geometry states this same idea; it is called the *side-angle-side postulate*.


*Side-Angle-Side postulate:* If two sides and the included angle of one triangle are congruent to the corresponding parts of another triangle, then the triangles are congruent.

Look at the following two triangles to see an example of this postulate:



**Practice**

Use the following figure to answer practice problems 16–20.



- |  |   |
|--|---|
| <p><b>16.</b> What kind of angles are <math>\angle ACB</math> and <math>\angle ECD</math>?</p> <p><b>17.</b> Is <math>\angle ACB \cong \angle ECD</math>?</p> <p><b>18.</b> <math>\overline{CE}</math> corresponds to _____.</p> | <p><b>19.</b> <math>\overline{BC}</math> corresponds to _____.</p> <p><b>20.</b> Is <math>\triangle ACB \cong \triangle EDC</math>?</p> |
|--|---|


**► Angle-Side-Angle (ASA) Postulate**

There is one more postulate that describes two congruent triangles. *Angle-side-angle* involves two angles and a side between them. The side is called an included side.

*Angle-Side-Angle postulate:* If two angles and the included side of one triangle are congruent to corresponding parts of another triangle, the triangles are congruent.

**CONGRUENT TRIANGLES**


Take a look at the following two triangles:



$$\triangle DEF \cong \triangle RST$$

**Practice**

Use the figure below to answer practice problems 21–25.



21.  $\angle TRS$  corresponds to  $\angle$ \_\_\_\_\_.

24. Is  $\triangle SRT \cong \triangle SQR$ ?


22.  $\angle QSR$  corresponds to  $\angle$ \_\_\_\_\_.

25. Is  $\angle T \cong \angle Q$ ?


23. Is  $\triangle RTS \cong \triangle RQS$ ?

State which postulate you would use to prove the two triangles congruent.


26.




27.




28.



29.




30.



**Skill Building until Next Time**

A carpenter's square is a tool used in carpentry. Similar tools are used for arts and crafts projects, as well as for wallpaper hanging. With the recent craze for home improvement, you may either have a carpenter's square or may be able to borrow one from a neighbor. A carpenter's square can be used to bisect an angle at the corner of a board. Get a carpenter's square and mark equal lengths  $\overline{WX}$  and  $\overline{WZ}$  along the edges. Put the carpenter's square on a board so that  $\overline{XY} = \overline{ZY}$ . Mark point Y and draw  $\overline{WY}$ .



- a) Which postulate can you use to show that  $\triangle WXY \cong \triangle WZY$ ?
- b) Why is  $\angle XWY \cong \angle ZWY$ ?





LESSON

# 8



# Triangles and the Pythagorean Theorem




## LESSON SUMMARY

In this lesson, you will learn the special names for the sides of a right triangle. You will also learn how to use the Pythagorean theorem to find missing parts of a right triangle and to determine whether three segments will make a right triangle.

**T**he right triangle is very important in geometry because it can be used in so many different ways. The Pythagorean theorem is just one of the special relationships that can be used to help solve problems and find missing information. Right triangles can be used to find solutions to problems that contain figures that are not even polygons.


### ► Parts of a Right Triangle

In a right triangle, the sides that meet to form the right angle are called the *legs*. The side opposite the right angle is called the *hypotenuse*. The hypotenuse is always the longest of the three sides. It is important that you can correctly identify the sides of a right triangle, regardless of what position the triangle is in.




**Practice**

$\triangle ABC$  is a right triangle.




1. Name the legs.
2. Name the hypotenuse.

$\triangle PQR$  is a right triangle.



3. Name the legs.
4. Name the hypotenuse.

$\triangle HLM$  is a right triangle.



5. Name the legs.
6. Name the hypotenuse.

## ► Review of Squares and Square Roots

Before you study the Pythagorean theorem, let's first review squares and square roots. Just like addition and subtraction are inverses, so are squares and square roots. In other words, they “undo” each other. To square a number, you multiply it by itself. A common mistake is to say that you multiply by two, since two is the exponent (the small raised number). But the exponent tells you how many times to use the base (bottom number) as a factor. For example,  $5^2$  means two factors of five, or five times five, which is twenty-five. Written algebraically, it looks like this:  $5^2 = 5 \times 5 = 25$ .

Twenty-five is a perfect square. It can be written as the product of two equal factors. It would be helpful for you to learn the first sixteen perfect squares. When completed, the following chart will be a useful reference. It is not necessarily important that you memorize the chart, but you need to understand how the numbers are generated. Even the most basic calculators can help you determine squares and square roots of larger numbers.

### Practice


7. Complete the chart.

NUMBER	SQUARE
1	1
2	4
3	9
4	
5	25
	36
7	49
8	
	81
10	100
11	
12	144
	169
14	196
15	
16	

## ► The Pythagorean Theorem

The Pythagorean theorem is one of the most famous theorems in mathematics. The Greek mathematician Pythagoras (circa 585–500 B.C.) is given credit for originating it. Evidence shows that it was used by the Egyptians and Babylonians for hundreds of years.


*Pythagorean theorem:* In a right triangle, the sum of the squares of the lengths of the legs is equal to the square of the length of the hypotenuse.




$$a^2 + b^2 = c^2$$

The Pythagorean theorem can be used to solve many real-life problems. Any unknown length can be found if you can make it a part of a right triangle. You only need to know two of the sides of a right triangle to find the third unknown side. A common mistake is always adding the squares of the two known lengths. You only add the squares of the legs when you are looking for the hypotenuse. If you know the hypotenuse and one of the legs, then you subtract the square of the leg from the square of the hypotenuse. Another common mistake is forgetting to take the square root as your final step. You just need to remember that you are not solving for the square of the side, but for the length of the side.

**Examples:** Find each missing length.



$$\begin{aligned} a^2 + b^2 &= c^2 \\ 12^2 + 9^2 &= c^2 \\ 144 + 81 &= c^2 \\ 225 &= c^2 \\ \sqrt{225} &= \sqrt{c^2} \\ 15 &= c \end{aligned}$$




$$\begin{aligned} a^2 + b^2 &= c^2 \\ 5^2 + b^2 &= 13^2 \\ 25 + b^2 &= 169 \\ b^2 &= 144 \\ \sqrt{b^2} &= \sqrt{144} \\ b &= 12 \end{aligned}$$


**Practice**

Find each missing length.


8.



9.



10.



11.  $a = 3, b = 4, c = \underline{\hspace{2cm}}$ .

12.  $a = \underline{\hspace{2cm}}, b = 9, c = 41$ .


13.  $a = 7, b = 24, c = \underline{\hspace{2cm}}$ .

► **Converse of the Pythagorean Theorem**

Another practical use of the Pythagorean theorem involves determining whether a triangle is acute, right, or obtuse. This involves using the converse of the Pythagorean theorem.

*Converse of the Pythagorean theorem:* If the square of the length of the longest side of a triangle is equal to the sum of the squares of the lengths of the two shorter sides, then the triangle is a right triangle.


**Examples:** Determine whether the following are right triangles. Note that  $c$  represents the length of the longest side in each triangle.



$$\begin{array}{c|c} c^2 & a^2 + b^2 \\ \hline 5^2 & 3^2 + 4^2 \\ 25 & 9 + 16 \end{array}$$

Since  $25 = 9 + 16$ , then the three sides with lengths of 5, 3, and 4 make a right triangle.

Here is another example:





$$\begin{array}{c|c} c^2 & a^2 + b^2 \\ \hline 6^2 & 4^2 + 5^2 \\ 36 & 16 + 25 \end{array}$$

Since  $36 \neq 16 + 25$ , then the three sides with lengths of 6, 4, and 5 do not make a right triangle.

**Practice**

The lengths of three sides of a triangle are given. Determine whether the triangles are right triangles.



\_\_\_\_\_ **16.** 25, 24, 7

\_\_\_\_\_ **17.** 5, 7, 9


\_\_\_\_\_ **18.** 15, 36, 39

\_\_\_\_\_ **19.** 9, 40, 41

### ► Acute and Obtuse Triangles


If you have determined that a triangle is not a right triangle, then you can determine whether it is an acute or obtuse triangle by using one of the following theorems:

*Theorem:* If the square of the length of the longest side is greater than the sum of the squares of the lengths of the other two shorter sides, then the triangle is obtuse.



$$c^2 > a^2 + b^2$$

*Theorem:* If the square of the length of the longest side is less than the sum of the squares of the lengths of the other two shorter sides, then the triangle is acute.



$$c^2 < a^2 + b^2$$



Let's take a look at a couple of examples to see these theorems in action.

**Example:** In  $\triangle ABC$ , the lengths of the sides are  $c = 24$ ,  $a = 15$ ,  $b = 20$ .

$$\begin{array}{r|l}
 c^2 & a^2 + b^2 \\
 \hline
 24^2 & 15^2 + 20^2 \\
 576 & 225 + 400 \\
 576 & 625
 \end{array}$$

Since  $576 < 625$ , this is an acute triangle.

**Example:** In  $\triangle ABC$ , the lengths of the sides are  $c = 9$ ,  $a = 5$ ,  $b = 7$ .

$$\begin{array}{r|l}
 c^2 & a^2 + b^2 \\
 \hline
 9 & 5^2 + 7^2 \\
 81 & 25 + 49 \\
 81 & 74
 \end{array}$$

Since  $81 > 74$ , this is an obtuse triangle.

**Practice**

The length of three sides of a triangle are given. Classify each triangle as acute, right, or obtuse.

**20.** 30, 40, 50

**21.** 10, 11, 13

**22.** 2, 10, 11


**23.** 7, 7, 10

**24.** 50, 14, 28

**25.** 5, 6, 7

**26.** 8, 12, 7

Use the following figure to answer question 27.



- 27.** How far up a building will an 18-foot ladder reach if the ladder's base is 5 feet from the building? Express your answer to the nearest foot. Solve the problem when the ladder is 3 feet from the building. Why would it be impractical to solve the problem if the base of the ladder was closer than 3 feet from the building?

### Skill Building until Next Time

Look around your home for examples of right triangles. How many can you find? Measure the three sides of the triangle to make sure it is a right triangle.



LESSON

9




# Properties of Polygons


## LESSON SUMMARY

In this lesson, you will learn how to determine if a figure is a polygon. You will also learn how to identify concave and convex polygons. You will learn how to classify polygons by their sides and how to find the measures of their interior and exterior angles.

**T**he word *polygon* comes from Greek words meaning “many angled.” A polygon is a closed plane figure formed by line segments. The line segments are called *sides* that intersect only at their endpoints, which are called *vertex points*.









Polygons



Not polygons

## PROPERTIES OF POLYGONS

A polygon is convex when no segment connecting two vertices (vertex points) contains points outside the polygon. In other words, if you wrapped a rubber band around a convex polygon, it would fit snugly without gaps. A concave polygon has at least one place that “caves in.”



Convex polygons

Concave polygons


### Practice

Classify each figure as a convex polygon, a concave polygon, or not a polygon.

1.




2.




3.




4.




5.




6.




7.



8.



9.




### ► Parts of a Polygon

Two sides of a polygon that intersect are called *consecutive* or *adjacent sides*. The endpoints of a side are called *consecutive* or *adjacent vertices*. The segment that connects two nonconsecutive vertices is called a *diagonal* of the polygon.

## ► Naming Polygons

When naming a polygon, you name its consecutive vertices in clockwise or counterclockwise order. Here are a few of the ways to name the following polygon:  $ABCDE$ ,  $DEABC$ , or  $EDCBA$ .



You cannot skip around when you name a polygon. For example, you cannot name the polygon  $BDEAC$ ,  $ECBAD$ , or  $ACEDB$ , to name just a few examples.


Polygons are classified by their number of sides.

NUMBER OF SIDES	POLYGON
3	triangle
4	quadrilateral
5	pentagon
6	hexagon
7	heptagon
8	octagon
9	nonagon
10	decagon


**Practice**

Name each polygon by the number of sides.


10.




13.




11.




14.



12.




15.





Which of the following are acceptable names for each polygon?



\_\_\_\_\_ 16. *DGFE*

\_\_\_\_\_ 19. *UPQRST*

\_\_\_\_\_ 17. *FDGE*

\_\_\_\_\_ 20. *RQSPUT*

\_\_\_\_\_ 18. *EFGD*


\_\_\_\_\_ 21. *UTSPQR*

### ► Finding the Measure of Interior Angles


There are two theorems that you can use to find the measure of interior angles of a convex polygon. One theorem works only for triangles. The other theorem works for all convex polygons, including triangles. Let's take a look at the theorem for triangles first.

*Theorem:* The sum of the interior angles of a triangle is  $180^\circ$ .


To illustrate this, cut a triangle from a piece of paper. Tear off the three angles or points of the triangle. Without overlapping the edges, put the vertex points together. They will form a straight line or straight angle. Remember that a straight angle is  $180^\circ$ ; therefore, the three angles of a triangle add up to  $180^\circ$ , as shown in the following figures.



You can find the sum of the interior angles of a convex polygon if you know how many sides the polygon has. Look at these figures. Do you see a pattern?



The diagram suggests that polygons can be divided into triangles. Since each triangle has  $180^\circ$ , multiply the number of triangles by 180 to get the sum of the interior angles.



Look for a pattern in the number of sides a polygon has and the number of triangles drawn from one vertex point. You will always have two fewer triangles than the number of sides of the polygon. You can write this as a general statement with the letter  $n$  representing the number of sides of the polygon.

*Theorem:* If a convex polygon has  $n$  sides, then its angle sum is given by this formula:  

$$S = 180(n - 2)$$

**Example:** Find the sum of the interior angles of a polygon with 12 sides.

**Solution:**  $n = 12$

$$S = 180(n - 2)$$

$$S = 180(12 - 2)$$

$$S = 180(10)$$

$$S = 1,800$$


Therefore, the sum of the interior angles of a 12-sided polygon is  $1,800^\circ$ .

### ► Finding the Measure of Exterior Angles

Use this theorem to find the measure of exterior angles of a convex polygon.

*Theorem:* The sum of the exterior angles of a convex polygon is always  $360^\circ$ .

To illustrate this theorem, picture yourself walking alongside a polygon. As you reach each vertex point, you will turn the number of degrees in the exterior angle. When you return to your starting point, you will have rotated  $360^\circ$ .




This figure shows this theorem using a pentagon. Do you see that this would be true for all polygons as stated in the theorem?

**Practice**

22. Complete the table for convex polygons

NUMBER OF SIDES	6	10	14	16
Interior $\angle$ Sum				
Exterior $\angle$ Sum				

23. The home plate used in baseball and softball has this shape:



As you can see, there are three right angles and two congruent obtuse angles. What are the measures of the two obtuse angles?

**Skill Building until Next Time**

Observe your surroundings today and see how many examples of convex and concave polygons you can find.

L E S S O N

10



# Quadrilaterals

## LESSON SUMMARY


In this lesson, you will learn how to name and classify special quadrilaterals. You will also learn how to use the special properties associated with parallelograms, rectangles, rhombuses, squares, and trapezoids.

**Q**uadrilaterals are one of the most commonly used figures in buildings, architecture, and design. The diagram on the next page shows the characteristics and relationships among the special quadrilaterals.

All four-sided polygons are classified as quadrilaterals. *Quadrilaterals* branch off in two distinctive branches: parallelograms and trapezoids. Trapezoids are quadrilaterals that have only one pair of opposite parallel sides. If the trapezoid has congruent legs, then the figure is an isosceles trapezoid. The diagram shows that an isosceles is one type of trapezoid, which is one type of quadrilateral. In other words, the figures become more specialized as the chart flows downward.

The other main branch of quadrilaterals consists of parallelograms. Parallelograms have two pairs of opposite parallel sides. Parallelograms branch off in two special categories: rectangles and rhombuses. A rectangle is a parallelogram with four congruent angles. A rhombus is a parallelogram with four congruent sides. A square is a parallelogram with four congruent angles and four congruent sides. In other words, a square is also a rectangle and a rhombus.

## QUADRILATERALS



### Practice

Use the diagram to determine whether each statement is true or false.


- |   |   |
|---|---|
| <p>_____ 1. All trapezoids are quadrilaterals.</p> <p>_____ 2. All parallelograms are rectangles.</p> <p>_____ 3. All rectangles are quadrilaterals.</p> <p>_____ 4. All parallelograms are squares.</p> <p>_____ 5. All squares are rhombuses.</p> | <p>_____ 6. All rectangles are parallelograms.</p> <p>_____ 7. All rhombuses are squares.</p> <p>_____ 8. All isosceles trapezoids are quadrilaterals.</p> <p>_____ 9. All squares are trapezoids.</p> <p>_____ 10. All quadrilaterals are squares.</p> |
|---|---|

## ► Properties of Parallelograms


The following properties of parallelograms will help you determine if a figure is a parallelogram or just a quadrilateral. The properties are also useful to determine measurements of angles, sides, and diagonals of parallelograms.

### Properties of Parallelograms

*Definition:* Opposite sides of a parallelogram are parallel.




*Theorem:* Opposite sides of a parallelogram are congruent.




*Theorem:* Opposite angles of a parallelogram are congruent.



*Theorem:* Consecutive angles of a parallelogram are supplementary.




*Theorem:* Diagonals of a parallelogram bisect each other.



## QUADRILATERALS


Note that diagonals of a parallelogram are not necessarily congruent. Watch out for this, because it is a common error.

**Examples:**  $BMAH$  is a parallelogram.



WE KNOW THAT	BECAUSE
$BM = 3$	Opposite sides are congruent.
$BH = 5$	Opposite sides are congruent.
$m\angle M = 135^\circ$	Opposite angles are congruent.
$m\angle A = 45^\circ$	Consecutive angles are supplementary.
$m\angle B = 45^\circ$	Opposite angles are congruent.


$WXYZ$  is a parallelogram.



WE KNOW THAT	BECAUSE
$m\angle XWZ = 90^\circ + 45^\circ = 135^\circ$	Angle-Addition-Postulate
$m\angle XYZ = 135^\circ$	Opposite angles are congruent.
$m\angle WXY = 45^\circ$	Consecutive angles are supplementary.
$m\angle WZY = 45^\circ$	Opposite angles are congruent.
$WO = 2$	Diagonals bisect each other.
$ZO = 6$	Diagonals bisect each other.

**Practice**

Use the following figure to find each side length and angle measure for parallelogram  $ABCD$  for questions 11–15.



\_\_\_\_\_ **11.**  $BC$


\_\_\_\_\_ **14.**  $m\angle A$

\_\_\_\_\_ **12.**  $DC$

\_\_\_\_\_ **15.**  $m\angle D$

\_\_\_\_\_ **13.**  $m\angle B$

Use the following figure to find each side length and angle measure for parallelogram  $PQRS$  for questions 16–20.



\_\_\_\_\_ **16.**  $SQ$

\_\_\_\_\_ **19.**  $m\angle SPQ$

\_\_\_\_\_ **17.**  $OR$

\_\_\_\_\_ **20.**  $m\angle SRQ$

\_\_\_\_\_ **18.**  $m\angle PQR$




### ► Other Special Properties

The rectangle, rhombus, and square have a few other special properties. First, remember that these figures are all parallelograms; therefore, they possess the same properties of any parallelogram. However, because these figures are special parallelograms, they also have additional special properties. Since a square is both a rectangle and a rhombus, a square possesses these same special properties.


*Theorem:* The diagonals of a rectangle are congruent.  
*Theorem:* The diagonals of a rhombus are perpendicular, and they bisect the angles of the rhombus.

**Examples:** *NEWS* is a rectangle.



WE KNOW THAT	BECAUSE
$XE = 6$	Diagonals bisect each other.
$NW = 12$	Diagonals of a rectangle are congruent.
$\angle NES = 20^\circ$	Alternate interior angles are congruent, when formed by parallel lines.
$\angle NSW = 90^\circ$	Definition of a rectangle.


ABCD is a rhombus.



WE KNOW THAT	BECAUSE
$m\angle BEC = 90^\circ$	Diagonals of a rhombus are perpendicular.
$m\angle DCE = 30^\circ$	Diagonals of a rhombus bisect the angles.
$m\angle BAD = 60^\circ$	Opposite angles are congruent.
$m\angle ABC = 120^\circ$	Consecutive angles are supplementary.
$m\angle ADC = 120^\circ$	Opposite angles are congruent.

**Practice**

Use the following figure to find the side length and angle measures for rectangle  $PQRS$  for questions 21–24.




\_\_\_\_\_ **21.**  $QS$

\_\_\_\_\_ **23.**  $m\angle QSR$

\_\_\_\_\_ **22.**  $OP$

\_\_\_\_\_ **24.**  $m\angle RPS$

Use the following figure to find the angle measures for rhombus  $GHJK$  for questions 25–30.



\_\_\_\_\_ **25.**  $m\angle MGK$

\_\_\_\_\_ **28.**  $m\angle GJK$


\_\_\_\_\_ **26.**  $m\angle GHJ$

\_\_\_\_\_ **29.**  $m\angle HJK$

\_\_\_\_\_ **27.**  $m\angle JMK$

\_\_\_\_\_ **30.**  $m\angle GKH$

Use the following diagram to answer question 31.



**31.** Explain how this Venn diagram and the flow chart shown at the beginning of this lesson show the relationships among quadrilaterals.

**Skill Building until Next Time**

Look around your environment to find examples of quadrilaterals and make a list of them. Then go down your list and see how many correct geometric names you can give for each item you named. For example, the state of Tennessee is a quadrilateral that is also a parallelogram. Your desktop may be a rectangle, a parallelogram, and a quadrilateral. The Venn diagram shown in practice problem 31 may be useful to you.

LESSON

# 11



# Ratio, Proportion, and Similarity

## LESSON SUMMARY


In this lesson, you will learn how to write and simplify ratios. You will also learn how to determine if two ratios are a proportion and how to use proportions to solve problems. In addition, you will learn how to determine if two triangles are similar.

**R**atios and proportions have many applications. Architects use them when they make scale models of buildings. Interior designers use scale drawings of rooms to decide furniture size and placement. Similar triangles can be used to find indirect measurements. Measurements of distances such as heights of tall buildings and the widths of large bodies of water can be found using similar triangles and proportions. Let's begin by looking at ratios.

### ► What Is a Ratio?

If you compare two quantities, then you have used a ratio. A *ratio* is the comparison of two numbers using division. The ratio of  $x$  to  $y$  can be written  $\frac{x}{y}$  or  $x:y$ . Ratios are usually expressed in simplest form.

**Examples:** Express each ratio in simplest form.



- (a)  $\frac{DR}{RY}$       (b)  $\frac{RY}{DR}$       (c)  $\frac{DR}{DY}$

**Solutions:**

- (a)  $\frac{7}{11}$       (b)  $\frac{11}{7}$       (c)  $\frac{7}{18}$

**Practice**

Express each ratio in simplest form.





- \_\_\_\_\_ 1.  $\frac{KB}{BT}$   
 \_\_\_\_\_ 2.  $\frac{BT}{KB}$   
 \_\_\_\_\_ 3.  $\frac{BT}{KT}$

**► What Is a Proportion?**

Since  $\frac{2}{4}$  and  $\frac{3}{6}$  are both equal to  $\frac{1}{2}$ , they are equal to each other. A statement that two ratios are equal is called a *proportion*. A proportion can be written in one of the following ways:

$$\frac{2}{4} = \frac{3}{6} \quad \text{or} \quad 2:4 = 3:6$$

The first and last numbers in a proportion are called the *extremes*. The middle numbers are called the *means*.



**Means-Extremes Property**

In a proportion, the product of the means equals the product of the extremes.

If  $\frac{a}{b} = \frac{c}{d}$ , then  $ad = bc$ .

If  $a:b = c:d$ , then  $ad = bc$ .

**Examples:** Tell whether each of the following is a proportion.

(a)  $\frac{3}{6} = \frac{1}{2}$

(b)  $2:5 = 4:10$

(c)  $\frac{1}{3} = \frac{2}{9}$

**Solutions:**

(a)  $3 \times 2 = 6 \times 1$

(b)  $2 \times 10 = 5 \times 4$

(c)  $1 \times 9 = 3 \times 2$

$6 = 6$

$20 = 20$

$9 \neq 6$

yes

yes

no

### Practice

State whether each of the following is a proportion.

\_\_\_\_\_ 4.  $\frac{2}{7} = \frac{4}{14}$

\_\_\_\_\_ 5.  $12:16 = 9:15$

\_\_\_\_\_ 6.  $2:3 = 3:2$

### ► Solving Proportion Problems

Proportions can also be used to solve problems. When three parts of a proportion are known, you can find the fourth part by using the *means-extremes property*.

**Examples:** Solve each proportion.

(a)  $\frac{4}{x} = \frac{2}{10}$

(b)  $\frac{a}{7} = \frac{12}{21}$

(c)  $\frac{5}{2} = \frac{15}{y}$

**Solutions:**

(a)  $2x = 4 \times 10$

(b)  $21a = 7 \times 12$

(c)  $5y = 2 \times 15$

$2x = 40$

$21a = 84$

$5y = 30$

$x = 20$

$a = 4$

$y = 6$

### Practice

Solve each proportion.

\_\_\_\_\_ 7.  $\frac{x}{8} = \frac{1}{2}$

\_\_\_\_\_ 10.  $\frac{2}{5} = \frac{a}{20}$

\_\_\_\_\_ 8.  $\frac{8}{y} = \frac{2}{11}$

\_\_\_\_\_ 11.  $\frac{8}{b} = \frac{4}{7}$

\_\_\_\_\_ 9.  $\frac{2}{7} = \frac{8}{z}$

\_\_\_\_\_ 12.  $\frac{9}{x} = \frac{36}{4}$

## ► Triangle Similarity

You can prove that two figures are similar by using the definition of *similar*. In other words, two figures are similar if you can show that the following two statements are true:

- (1) Corresponding angles are congruent.
- (2) Corresponding sides are in proportion.


In addition to using the definition of similar, you can use three other methods for proving that two triangles are similar. The three methods are called the angle-angle postulate, the side-side-side postulate, and the side-angle-side postulate.

If you know the measurements of two angles of a triangle, can you find the measurement of the third angle? Yes, from Lesson 9, you know that the sum of the three angles of a triangle is  $180^\circ$ . Therefore, if two angles of one triangle are congruent to two angles of another triangle, then their third angles must also be congruent. This will help you understand the next postulate. You should know that the symbol used for similarity is  $\sim$ .


*Angle-Angle postulate (AA postulate):* If two angles of one triangle are congruent to two angles of another triangle, then the triangles are similar.

**Examples:** Are these triangles similar?

(a)



(b)



**Solutions:**


- (a)  $\angle A \cong \angle D$ , given  
 $\angle BCA \cong \angle ECD$ , vertical  $\angle$ 's  
 are  $\cong$   
 $\triangle ABC \sim \triangle DEC$ , AA postulate

- (b)  $m\angle J = 180 - (60 + 40)$   
 $m\angle J = 80$   
 $\triangle AJT \sim \triangle YZX$ , AA postulate


**Practice**

State whether the triangles are similar.


\_\_\_\_\_ **13.**




\_\_\_\_\_ **14.**



\_\_\_\_\_ **15.**



\_\_\_\_\_ **16.**





## RATIO, PROPORTION, AND SIMILARITY


Here are two more postulates you can use to prove that two triangles are similar:

*Side-Side-Side postulate (SSS postulate):* If the lengths of the corresponding sides of two triangles are proportional, then the triangles are similar.

*Side-Angle-Side postulate (SAS postulate):* If the lengths of two pairs of corresponding sides of two triangles are proportional and the corresponding included angles are congruent, then the triangles are similar.

**Examples:** Which postulate, if any, could you use to prove that the triangles are similar?


(a)



$$\frac{3}{4} \stackrel{?}{=} \frac{12}{16}$$
$$3 \times 16 = 4 \times 12$$
$$48 = 48$$

SAS postulate

(b)




$$\frac{6}{15} \stackrel{?}{=} \frac{10}{25}$$
$$6 \times 25 = 15 \times 10$$
$$150 = 150$$

SSS postulate

$$\frac{6}{15} \stackrel{?}{=} \frac{8}{20}$$
$$6 \times 20 = 15 \times 8$$
$$120 = 120$$
$$\frac{8}{20} \stackrel{?}{=} \frac{10}{25}$$
$$8 \times 25 = 20 \times 10$$
$$200 = 200$$

(c)




$$\frac{3}{4} \stackrel{?}{=} \frac{9}{10}$$

$$3 \times 10 = 4 \times 9$$

$$30 \neq 36$$

none

(d)



$$\frac{2}{3} \stackrel{?}{=} \frac{4}{6}$$

$$2 \times 6 = 3 \times 4$$


$12 = 12$ , but the included  $\angle$ 's are not necessarily  $\cong$

none


### Practice

Which postulate, if any, could you use to prove that the triangles are similar?


17.




18.




19.




20.




21.




22.



23.



24.



### Skill Building until Next Time

Outside your school, find two objects that are fairly close to each other. One should be short enough to measure with a measuring tape. Some good choices would be a stop sign or a fireplug. The other should be too tall to measure with a measuring tape. A flag pole or a light post would be good choices. Measure the length of the shadows of the short and the tall objects. Set up a proportion:

$$\frac{\text{height of the flag pole}}{\text{shadow of the flag pole}} = \frac{\text{height of the stop sign}}{\text{shadow of the stop sign}}$$

Replace the height of the flag pole with  $x$  and the other parts of the proportion with the measurements that you found. Solve by using the *means-extremes property*.



LESSON

# 12



# Perimeter of Polygons


## LESSON SUMMARY

In this lesson, you will learn how to find the distance around convex and concave polygons. You will also learn how to use formulas to find the perimeter of polygons.


**P**erimeter is found by measuring the distance around an object. Crown molding around a room and a fence around a garden are just two examples of where you might need to find a perimeter. You simply add all the lengths of the sides of an irregular shape to find its perimeter. You can find the perimeters of polygons, which have a uniform shape by using formulas. Carpenters and landscape architects use perimeters on a regular basis.

**Examples:** Find the perimeter of each polygon.

(a)




(b)



**Solutions:**

(a)  $\text{perimeter} = 1 + 4 + 2 + 5 + 6$   
 $= 18 \text{ cm}$

(b)




$\text{perimeter} = 7 + 7 + 5 + 4 + 2 + 3$   
 $= 28$


**Practice**

Find the perimeter of each polygon.


\_\_\_\_\_ **1.**




\_\_\_\_\_ **2.**



\_\_\_\_\_ **3.**




\_\_\_\_\_ **4.**



### ► Perimeter Formulas


For certain shapes, you can find the perimeter in a more efficient manner. Using a standard formula for a particular shape is faster and easier than adding all the sides. Here are the most commonly used perimeter formulas:

*Square*



$$p = 4s$$

*Rectangle*




$$p = 2l + 2w$$

or

$$p = 2(l + w)$$


*Regular pentagon*



$$p = 5s$$




**Examples:** Use a formula to find each perimeter.



$$p = 4s$$

$$p = 4(8)$$

$$p = 32 \text{ cm}$$




$$p = 2l + 2w \quad \text{or} \quad p = 2(l + w)$$

$$p = 2(7) + 2(3) \quad p = 2(7 + 3)$$

$$p = 14 + 6 \quad p = 2(10)$$

$$p = 20 \text{ cm} \quad p = 20 \text{ cm}$$

Either of these formulas will always work.  
You can choose which one is easier for you.



$$p = 5s$$

$$p = 5(4)$$


$$p = 20 \text{ cm}$$

A *regular polygon* is a polygon with all angles congruent and all sides congruent. So if you know the length of one side of a regular polygon, all you need to do to find its perimeter is multiply the number of sides by the length of a side. In other words,  $p = ns$ , where  $n$  = number of sides and  $s$  = the length of each side.


**Practice**

Use a formula to find the perimeter of each polygon.


\_\_\_\_\_ **5.**




\_\_\_\_\_ **7.**



\_\_\_\_\_ **6.**



\_\_\_\_\_ **8.**



\_\_\_\_\_ **9.** square:  $s = 20$  cm

\_\_\_\_\_ **10.** rectangle:  $l = 12$  in.,  $w = 6$  in.

\_\_\_\_\_ **11.** regular octagon:  $s = 10$  in.


\_\_\_\_\_ **12.** regular nonagon:  $s = 7$  ft.

### ► Working Backward

You can also use a perimeter formula to find the length of one side of a polygon if you know the perimeter and work backward.

**Examples:**

(a)




$$p = 20 \text{ cm}$$

$$s = \underline{\hspace{2cm}}$$

**Solutions:**

$$\begin{aligned} \text{(a)} \quad p &= 4s \\ 20 &= 4s \\ s &= 5 \text{ cm} \end{aligned}$$

(b)



$$p = 32$$

$$w = \underline{\hspace{2cm}}$$

$$l = \underline{\hspace{2cm}}$$

$$\begin{aligned} \text{(b)} \quad w &= 5 \\ p &= 2l + 2w \\ 32 &= 2l + 2(5) \\ 32 &= 2l + 10 \\ 22 &= 2l \\ l &= 11 \end{aligned}$$

(c) regular hexagon


$$p = 24 \text{ mm}$$

$$\begin{aligned} \text{(c)} \quad p &= 6s \\ 24 &= 6s \\ s &= 4 \text{ mm} \end{aligned}$$

**Practice**

Find the length of the indicated side(s).

**13.**




$$p = 12 \text{ m}$$

$$w = \underline{\hspace{2cm}}$$

$$l = \underline{\hspace{2cm}}$$

**14.**




$$p = 30 \text{ cm}$$

$$w = \underline{\hspace{2cm}}$$

$$l = \underline{\hspace{2cm}}$$


**15.**



$$p = 35 \text{ in.}$$

$$s = \underline{\hspace{2cm}}$$

**16.**



$$p = 32 \text{ yd.}$$

$$s = \underline{\hspace{2cm}}$$

**17.** equilateral triangle:  $p = 60 \text{ cm}$   $s = \underline{\hspace{2cm}}$

**18.** regular octagon:  $p = 80 \text{ cm}$   $s = \underline{\hspace{2cm}}$

**19.** regular hexagon:  $p = 48 \text{ cm}$   $s = \underline{\hspace{2cm}}$

**20.** regular decagon:  $p = 230 \text{ cm}$   $s = \underline{\hspace{2cm}}$

Find answers to these practical problems.

- 21.** Suppose you want to frame an 8-inch by 10-inch picture. How much molding will you need to buy?
- 22.** A fence must be placed around your vegetable garden. The dimensions of the garden are 30 feet by 10 feet. How much fencing will you need?
- 23.** You have a square dining room that you would like to trim with crown molding. If the length of the room is 17 feet, how much crown molding should you purchase?
- 24.** Suppose you want to hang Christmas icicle lights around your house. If your house is 32 feet by 40 feet and one package of lights is 9 feet long, how many packages of lights should you buy?

### Skill Building until Next Time

Use a tape measure to find the perimeter of your room, the door of your room, and a window in your room. Use formulas when possible.

LESSON

13



# Area of Polygons

## LESSON SUMMARY

In this lesson, you will learn how to use formulas to find the area of rectangles, parallelograms, triangles, and trapezoids. You will also learn how to use the formulas to work backward to find missing lengths of polygons.

**P**eople often confuse area and perimeter. As you learned in Lesson 12, perimeter is the distance around an object. In this lesson, you'll work with *area*, which is the amount of surface covered by an object. For example, the number of tiles on a kitchen floor would be found by using an area formula, while the amount of baseboard used to surround the room would be found by using a perimeter formula. Perimeter is always expressed in linear units. Area is always expressed in square units. It should be obvious that you could not measure the amount of surface that is covered by an object by simply measuring it in one direction. You need to measure the object in two directions, like a square. A flat surface has two dimensions: length and width. When you multiply a number by itself, the number is said to be squared. In the same way, when two units of measurement are multiplied by each other, as in area, the unit is expressed in square units. By looking at a tiled floor, it is easy to see that *area* refers to how many squares it takes to cover a surface.

## ► Finding the Area of a Rectangle

For a rectangle, the base can be any side. The base length is represented by  $b$ . The sides perpendicular to the base are referred to as the height. The height is referred to as  $h$ . The base is often called the length,  $l$ , and the height is often called the width,  $w$ . Length,  $l$ , and width,  $w$ , are used in the same manner as base,  $b$ , and height,  $h$ . This book uses *base* and *height*.


Here is a useful theorem you can use to find the area of a rectangle.

*Theorem:* The area ( $A$ ) of a rectangle is the product of its base length ( $b$ ) and its height ( $h$ ).

$$A = bh$$

**Examples:** Find the area of each rectangle.

(a)



5 cm

$$A = bh$$

$$A = 5(8)$$

$$A = 40 \text{ cm}^2$$

(b)



10 ft.





$$A = bh$$

$$A = 10(2)$$

$$A = 20 \text{ ft.}^2$$

**Practice**

Find the area of each rectangle.



► **Finding the Length of an Unknown Side of a Rectangle**

You can also use the area formula for a rectangle to find the length of an unknown side if you know the area.

**Examples:** For each rectangle, find the length of the indicated sides.

(a)

**Solution:**  $A = bh$

$$32 = 8h$$

$$h = 4 \text{ cm}$$




$$A = 32 \text{ cm}^2$$

$$b = \underline{\hspace{2cm}}$$

$$h = \underline{\hspace{2cm}}$$






**Solution:**  $A = bh$   
 $84 = b(12)$   
 $b = 7 \text{ ft}$

$A = 84 \text{ ft.}^2$   
 $b = \underline{\hspace{2cm}}$   
 $h = \underline{\hspace{2cm}}$

**Practice**


For each figure, find the length of the indicated side(s).



$A = 56 \text{ in.}^2$   
 $b = \underline{\hspace{2cm}}$   
 $h = \underline{\hspace{2cm}}$

**7.**  $A = 25 \text{ yd.}^2$   
 $b = 4 \text{ yd.}$   
 $h = \underline{\hspace{2cm}}$

**8.**  $A = 150 \text{ cm}^2$   
 $b = \underline{\hspace{2cm}}$   
 $h = 10 \text{ cm}$





$A = 75 \text{ m}^2$   
 $b = \underline{\hspace{2cm}}$   
 $h = \underline{\hspace{2cm}}$


► **Finding the Area and Unknown Sides of a Parallelogram**

Any side of a parallelogram can be called the base. The height is the length of the altitude. The altitude is a segment perpendicular to the base.

AREA OF POLYGONS



Draw an altitude of a parallelogram. Cut along the altitude to separate the parallelogram into two pieces. Fit the two pieces together to form a rectangle. You'll find that the base and height of the rectangle coincide with the base and altitude of the parallelogram.




Using this information, can you predict the area formula of a parallelogram? Take a moment to make your prediction, then look at the following theorem.

*Theorem:* The area ( $A$ ) of a parallelogram is the product of its base length ( $b$ ) and its height ( $h$ ).


$$A = bh$$

**Examples:** Find the area of the parallelogram.



**Solution:**  $A = bh$   
 $A = 12(9)$   
 $A = 108 \text{ in.}^2$

Find the indicated lengths.




$A = 40 \text{ cm}^2$       **Solution:**  $A = bh$   
 $b = \underline{\hspace{2cm}}$                        $40 = 10h$   
 $h = \underline{\hspace{2cm}}$                        $h = 4 \text{ cm}$

Note that 5 cm is unnecessary information for this problem. Recall that the base and height must be perpendicular to each other.

**Practice**

Find the indicated information.

9.




$A = \underline{\hspace{2cm}}$

$b = \underline{\hspace{2cm}}$

$h = \underline{\hspace{2cm}}$

10.




$A = \underline{\hspace{2cm}}$

$b = \underline{\hspace{2cm}}$

$h = \underline{\hspace{2cm}}$

11.




$A = 60 \text{ cm}^2$

$b = \underline{\hspace{2cm}}$

$h = \underline{\hspace{2cm}}$

12.




$A = 28 \text{ ft.}^2$

$b = \underline{\hspace{2cm}}$

$h = \underline{\hspace{2cm}}$

**► Finding the Area and Unknown Sides of a Triangle**

Look at the following figures and try to predict the area formula for a triangle.



Now see if your prediction is correct by reading the theorem on the next page.


**AREA OF POLYGONS**

*Theorem:* The area ( $A$ ) of any triangle is half the product of its base length ( $b$ ) and height ( $h$ ).

$$A = \frac{1}{2}bh$$

**Examples:** Find the area.

(a)




$$A = \frac{1}{2}bh$$

$$A = \frac{1}{2}(10)(6)$$

$$A = 30 \text{ in.}^2$$

(b)



$$A = \frac{1}{2}bh$$


$$A = \frac{1}{2}(7)(4)$$

$$A = 14 \text{ cm}^2$$

**Practice**

Find each area.


\_\_\_\_\_ **13.**



\_\_\_\_\_ **15.** Triangle:  $b = 10$  feet,  $h = 7$  feet


\_\_\_\_\_ **16.** Triangle:  $b = 13$  inches,  $h = 26$  inches

\_\_\_\_\_ **14.**



Find the indicated length.


17.



$$A = 77 \text{ cm}^2$$

$$h = \underline{\hspace{2cm}}$$


19.



$$A = 51 \text{ in.}^2$$

$$b = \underline{\hspace{2cm}}$$

18.




$$A = 68 \text{ ft.}^2$$

$$h = \underline{\hspace{2cm}}$$

► **Finding the Area and Unknown Sides of a Trapezoid**

Can you predict the area formula for a trapezoid? Look at these figures:



Area of the trapezoid = area of two triangles

$$= \frac{1}{2}b_1h + \frac{1}{2}b_2h$$


$$= \frac{1}{2}h(b_1 + b_2)$$

*Theorem:* The area of a trapezoid is half the product of the height and the sum of the base lengths ( $b_1 + b_2$ ).

$$A = \frac{1}{2}h(b_1 + b_2)$$

**Examples:** Find the area or indicated length.

(a)




$$A = \frac{1}{2}h(b_1 + b_2)$$

$$A = \frac{1}{2}(6)(7 + 12)$$

$$A = \frac{1}{2}(6)(19)$$

$$A = 57 \text{ cm}^2$$

(b)



$$A = \frac{1}{2}h(b_1 + b_2)$$

$$130 = \frac{1}{2}(10)(b_1 + 16)$$

$$130 = 5(b_1 + 16)$$

$$130 = 5b_1 + 80$$


$$50 = 5b_1$$

$$b_1 = 10 \text{ cm}$$

**Practice**


Find the area or indicated length.

20.




$A = \underline{\hspace{2cm}}$

21.




$A = 90 \text{ cm}^2$   
 $b_1 = \underline{\hspace{2cm}}$

22.



$A = \underline{\hspace{2cm}}$

23.



$A = 75 \text{ in.}^2$

$h = \underline{\hspace{2cm}}$

24.  $h = 6 \text{ cm}$

$b_1 = 5 \text{ cm}$

$b_2 = 10 \text{ cm}$

$A = \underline{\hspace{2cm}}$

25.  $A = 92 \text{ m}^2$

$h = 8 \text{ m}$

$b_1 = 14 \text{ m}$

$b_2 = \underline{\hspace{2cm}}$

### Skill Building until Next Time

Try to find an object shaped like each of the polygons in this lesson. Examples would be a rectangular window pane, a square floor tile, a table leg shaped like a trapezoid, a quilt piece shaped like a parallelogram, and a triangular scarf. Measure and find the area of each.

LESSON

14



# Surface Area of Prisms

## LESSON SUMMARY


In this lesson, you will learn how to identify various parts of a prism and how to use formulas to find the surface area of a prism.

**A** right rectangular prism is a shape you see every day. Examples are bricks, cereal boxes, and some buildings. Sometimes, it is necessary to completely cover these objects. You might need to wrap a package for Christmas, determine how much siding is needed to cover a house, or make a label for a box.

A rectangular prism has six faces. The faces are parts of planes that form sides of solid figures. A rectangular prism has twelve edges. The edges are the segments formed by the intersection of two faces of a solid figure. A rectangular prism has eight vertices. Vertices are the points where the edges meet.



**Example:**




$ABCD$  is a face

$CG$  is an edge

$G$  is a vertex

Edges that are parallel to each other have the same measurements.

**Example:**




$PQ = 2$  cm

$LO = 3$  cm

$LP = 6$  cm

**Practice**

State the length of each side.



1.  $HL = \underline{\hspace{2cm}}$

6.  $LM = \underline{\hspace{2cm}}$

2.  $KL = \underline{\hspace{2cm}}$

7. How many edges does a rectangular solid have?

3.  $GJ = \underline{\hspace{2cm}}$

8. How many faces does a rectangular solid have?



4.  $GK = \underline{\hspace{2cm}}$

9. How many vertices does a rectangular solid have?

5.  $JL = \underline{\hspace{2cm}}$

**► Using a Formula to Find the Surface Area of a Rectangular Prism**

You can take a box apart, fold out all the sides, and you will have six rectangles. You will have three pairs that are the same size. You could find the area of each rectangle and add up all their areas to obtain the total area.




You could make a chart to help you organize all the faces, or you could use a formula.


## SURFACE AREA OF PRISMS

*Theorem:* The surface area (S.A.) of a rectangular prism is twice the sum of the length ( $l$ ) times the width ( $w$ ), the width ( $w$ ) times the height ( $h$ ), and the length ( $l$ ) times the height ( $h$ ).

$$S.A. = 2(lw + wh + lh)$$



**Example:** Find the surface area of the rectangular prism.



$$l = 3 \text{ cm}, w = 2 \text{ cm}, h = 5 \text{ cm}$$

$$S.A. = 2(lw + wh + lh)$$

$$S.A. = 2[(3)(2) + (2)(5) + (3)(5)]$$





$$S.A. = 2(6 + 10 + 15)$$

$$S.A. = 2(31)$$

$$S.A. = 62 \text{ cm}^2$$

**Practice**

Find the surface area of each rectangular prism.




\_\_\_\_\_ **14.**  $l = 9 \text{ in.}, w = 3 \text{ in.}, h = 5 \text{ in.}$

\_\_\_\_\_ **15.**  $l = 5 \text{ ft.}, w = 10 \text{ ft.}, h = 4 \text{ ft.}$

**The Surface Area of a Cube**


A cube is a special rectangular prism. All the edges of a cube have the same length. All six faces have the same area.



*Theorem:* The surface area of a cube is six times the edge ( $e$ ) squared.

$$\text{S.A.} = 6e^2$$

**Example:** Find the surface area of the cube.




$$\begin{aligned} \text{S.A.} &= 6e^2 \\ \text{S.A.} &= 6(3)^2 \\ \text{S.A.} &= 6(9) \\ \text{S.A.} &= 54 \text{ cm}^2 \end{aligned}$$


**Practice**

Find the surface area of each cube.


\_\_\_\_\_ **16.**




\_\_\_\_\_ **18.**



\_\_\_\_\_ **17.**



\_\_\_\_\_ **19.**



\_\_\_\_\_ **20.**  $e = 5.8 \text{ cm}$

\_\_\_\_\_ **21.** The surface area of a cube is  $486 \text{ ft.}^2$   
Find the length of each edge.

**Skill Building until Next Time**

Take a juice box and measure its length, width, and height. Use the formula to find its surface area. Now take a break and drink the juice. Unfold the empty juice box. Find the area of the rectangles. Compare the two answers.

LESSON

15



# Volume of Prisms and Pyramids

## LESSON SUMMARY

In this lesson, you will learn how to find the volume of prisms and pyramids using formulas.

**W**hen you are interested in finding out how much a refrigerator holds or the amount of storage space in a closet, you are looking for the volume of a prism. A *prism* is a polyhedron with a pair of parallel bases. A *polyhedron* is a three-dimensional figure with all surfaces that are polygons. Volume is expressed in cubic units. Just like an ice tray is filled with cubes of ice, volume tells you how many cubic units will fit into a space.

### ► Volume of Prisms

Prisms can have bases in the shape of any polygon. A prism with a rectangle for its bases is called a rectangular prism. A prism with triangles for its bases is referred to as a triangular prism. A prism with hexagons as its bases is called a hexagonal prism, and so on. Prisms can be right or oblique. A right prism is a prism with its bases perpendicular to its sides, meaning they form right angles. Bases of oblique prisms do not form right angles with their sides. An example of an oblique prism is the Leaning Tower of Pisa. In this lesson, you will concentrate on the volume of right prisms. When we refer to prisms in this lesson, you can assume we mean a right prism.

*Theorem:* To find the volume ( $V$ ) of a rectangular prism, multiply the length ( $l$ ) by the width ( $w$ ) and by the height ( $h$ ).

$$V = lwh$$


The volume of a rectangular prism could also be stated in another way. The area of the base of a rectangular prism is the length times the width, the same length and width used with the height to find the volume. In other words, the area of the base ( $B$ ) times the height. This same approach can also be applied to other solid figures. In Lesson 13, you studied ways to find the area of most polygons, but not all. However, if you are given the area of the base and the height of the figure, then it is possible to find its volume. Likewise, if you do know how to find the area of the base ( $B$ ), use the formula for that shape, then multiply by the height ( $h$ ) of the figure.

*Theorem:* To find the volume ( $V$ ) of any prism, multiply the area of a base ( $B$ ) by the height ( $h$ ).

$$V = Bh$$

**Examples:** Find the volume of each prism.

(a)




$$V = lwh$$

$$V = (12)(6)(6)$$

$$V = 432 \text{ in.}^3$$

(b)



**Solution:** First, find the area of a base. As you remember from Lesson 13, the formula for finding the area of any triangle is  $A = \frac{1}{2}bh$ .

$$A = \frac{1}{2}(3)(4)$$

$$A = \frac{1}{2}(12)$$

$$A = 6 \text{ in.}^2$$


The area,  $A$ , becomes the base,  $B$ , in the volume formula  $V = Bh$ .

$$V = Bh$$

$$V = (6)(6)$$

$$V = 36 \text{ in.}^3$$

(c)



$$B = 30 \text{ ft.}^2 \quad \textbf{Solution: } V = Bh$$

$$h = 5 \text{ ft.}$$

$$V = (30)(5)$$

$$V = 150 \text{ ft.}^3$$

(d) Prism:  $B = 42 \text{ m}^2$ ,  $h = 10 \text{ m}$  **Solution:**  $V = Bh$

$$V = (42)(10)$$


$$V = 420 \text{ m}^3$$




**Practice**

Find the volume of each prism.


\_\_\_\_\_ **1.**




\_\_\_\_\_ **2.**




\_\_\_\_\_ **3.**




\_\_\_\_\_ **4.**




\_\_\_\_\_ **5.**




\_\_\_\_\_ **6.**




\_\_\_\_\_ **7.**



\_\_\_\_\_ **8.**



\_\_\_\_\_ **9.**



\_\_\_\_\_ **10.** Prism:  $B = 80 \text{ cm}^2$ ,  $h = 14 \text{ cm}$

\_\_\_\_\_ **11.** Prism:  $B = 48 \text{ ft.}^2$ ,  $h = 12 \text{ ft.}$

\_\_\_\_\_ **12.** Prism:  $B = 85 \text{ m}^2$ ,  $h = 8 \text{ m}$


## ► Volume of a Cube

Recall that the edges ( $e$ ) of a cube all have the same measurement; therefore, if you replace the length ( $l$ ), width ( $w$ ), and height ( $h$ ) with the measurement of the edge of the cube, then you will have the formula for the volume of the cube,  $V = e^3$ .

*Theorem:* The volume of a cube is determined by cubing the length of the edge.

$$V = e^3$$

**Example:** Find the volume of the cube.



$$V = e^3$$


$$V = (5)^3$$

$$V = 125 \text{ m}^3$$

### Practice

Find the volume of each cube.


\_\_\_\_\_ **13.**



\_\_\_\_\_ **15.** Cube:  $e = 4$  in.


\_\_\_\_\_ **16.** Cube:  $e = 9$  cm

\_\_\_\_\_ **14.**




## ► Volume of a Pyramid


A regular pyramid is a polyhedron with a base that is a regular polygon and a vertex point that lies directly over the center of the base.



If you have a pyramid and a rectangular prism with the same length, width, and height, you would find that it would take three of the pyramids to fill the prism. In other words, one-third of the volume of the prism is the volume of the pyramid.




$$V = lwh \text{ or } V = Bh$$



$$V = \frac{1}{3}lwh \text{ or } V = \frac{1}{3}Bh$$

**Examples:** Find the volume of each pyramid.

(a)




$$V = \frac{1}{3}lwh$$

$$V = \frac{1}{3}(5)(3)(6)$$

$$V = 30 \text{ cm}^3$$

(b)



$$V = \frac{1}{3}Bh$$

$$V = \frac{1}{3}(50)(9)$$

$$V = 150 \text{ in.}^3$$

(c) Regular pyramid:  $B = 36 \text{ ft.}^2$ ,  $h = 8 \text{ ft.}$

**Solution:**  $V = \frac{1}{3}Bh$


$$V = \frac{1}{3}(36)(8)$$

$$V = 96 \text{ ft.}^3$$


**Practice**

Find the volume of each pyramid.


\_\_\_\_\_ **17.**




\_\_\_\_\_ **20.**




\_\_\_\_\_ **18.**




\_\_\_\_\_ **21.**



\_\_\_\_\_ **19.**



\_\_\_\_\_ **22.**



\_\_\_\_\_ **23.** Regular pyramid:  $B = 64$  in.<sup>2</sup>,  $h = 6$  in.

\_\_\_\_\_ **24.** Regular pyramid:  $B = 156$  ft.<sup>2</sup>,  $h = 8$  ft.

\_\_\_\_\_ **25.** Regular pyramid:  $B = 3$  cm<sup>2</sup>,  $h = 1$  cm

**Skill Building until Next Time**

Take a box of saltine crackers, the kind with four packages of crackers inside. Measure the length, width, and height of the cracker box and find its volume. Now measure the length, width, and height of the stick of crackers and find its volume. Compare the two volumes.

LESSON

# 16



## Working with Circles and Circular Figures

### LESSON SUMMARY

In this lesson, you will learn about the irrational number,  $\pi$ . You will also learn to use formulas to find the circumference and area of circles, the surface area and volume of cylinders and spheres, and the volume of cones.

**B**efore you begin to work with circles and circular figures, you need to know about the irrational number,  $\pi$  (pronounced “pie”). Over 2,000 years ago, mathematicians approximated the value of the ratio of the distance around a circle to the distance across a circle to be approximately 3. Years later, this value was named with the Greek letter  $\pi$ . The exact value of  $\pi$  is still a mathematical mystery.  $\pi$  is an irrational number. A *rational number* is a number that can be written as a ratio, a fraction, or a terminating or repeating decimal.  $\pi$  has been computed in various ways over the past several hundred years, but no one has been able to find a decimal value of  $\pi$  where the decimal terminates or develops a repeating pattern. With the recent progress in computers, the value of  $\pi$  has been computed to over fifty billion decimal places, but there is still no termination or repeating group of digits.

$$\pi = \frac{\text{circumference}}{\text{diameter}}$$

The most commonly used approximations for  $\pi$  are  $\frac{22}{7}$  and 3.14. These are not the values of  $\pi$ , only rounded approximations. You may have a  $\pi$  key on your calculator. This key will give you an approximation for  $\pi$  that varies according to how many digits your calculator holds.

## ► Circumference of a Circle

Now that you know about the irrational number  $\pi$ , it's time to start working with circles. The distance around a circle is called its *circumference*. There are many reasons why people need to find a circle's circumference. For example, the amount of lace edge around a circular skirt can be found by using the circumference formula. The amount of fencing for a circular garden is another example of when you need the circumference formula.

Since  $\pi$  is the ratio of circumference to diameter, then the approximation of  $\pi$  times the diameter of the circle gives you the circumference of the circle. The diameter of a circle is the distance across a circle through its center. A radius is the distance from the center to the edge. One-half the diameter is equal to the radius or two radii are equal to the length of the diameter.

$$d = 2r \text{ or } r = \frac{1}{2}d$$


Here is a theorem that will help you solve circumference problems:

*Theorem:* The circumference of any circle is the product of its diameter and  $\pi$ .  
 $C = \pi d$  or  $C = 2\pi r$

Since  $\pi$  is approximately 3.14 (and not exactly equal to 3.14), after you substitute the value 3.14 for  $\pi$  in the formula, you should use  $\approx$  instead of  $=$ . The symbol  $\approx$  means *approximately equal to*.

**Examples:** Find the approximate circumference of each circle. Use the approximation of 3.14 for  $\pi$ .


(a)



**Solution:** Use the  $C = \pi d$  formula, since the diameter of the circle is given.

$$\begin{aligned} C &= \pi d \\ C &\approx (3.14)(10) \\ C &\approx 31.4 \text{ cm} \end{aligned}$$

(b)



**Solution:** Use the  $C = 2\pi r$  formula, since the radius of the circle is given.


$$\begin{aligned} C &= 2\pi r \\ C &\approx 2(3.14)(5) \\ C &\approx 31.4 \text{ cm} \end{aligned}$$

Notice that these two circles have the same circumference because a circle with a diameter of 10 cm has a radius of 5 cm. You pick which formula to use based on what information you are given—either the circle’s radius or its diameter.

**Practice**

Find the approximate circumference of each circle shown or described. Use 3.14 for  $\pi$ .


\_\_\_\_\_ 1.



\_\_\_\_\_ 3.  $d = 7$  m


\_\_\_\_\_ 4.  $r = 25$  m

\_\_\_\_\_ 2.



► **Area of a Circle**

To understand the area of a circle, take a look at the following figure. Imagine a circle that is cut into wedges and rearranged to form a shape that resembles a parallelogram.




*Theorem:* The area ( $A$ ) of a circle is the product of  $\pi$  and the square of the radius ( $r$ ).  
 $A = \pi r^2$

Notice that this formula squares the radius, not the diameter, so if you are given the diameter, you should divide it by two or multiply it by one-half to obtain the radius. Some people will mistakenly think that squaring the radius and doubling it to get the diameter are the same. They are the same only when the radius is two. Otherwise, two times a number and the square of a number are very different.



**Examples:** Find the approximate area for each circle. Use 3.14 for  $\pi$ .

(a)




$$A = \pi r^2$$

$$A \approx (3.14)(6)^2$$

$$A \approx (3.14)(36)$$

$$A \approx 113.04 \text{ cm}^2$$

(b)



$$r = \frac{1}{2}d$$

$$r = \frac{1}{2}(14)$$

$$r = 7 \text{ in.}$$

$$A = \pi r^2$$

$$A \approx (3.14)(7)^2$$


$$A \approx (3.14)(49)$$

$$A \approx 153.86 \text{ in.}^2$$

**Practice**

Find the approximate area for each circle shown or described. Use 3.14 for  $\pi$ .


\_\_\_\_\_ **5.**



\_\_\_\_\_ **7.**  $r = 13 \text{ cm}$


\_\_\_\_\_ **8.**  $d = 24 \text{ in.}$

\_\_\_\_\_ **6.**



### ► Surface Area of a Cylinder

When you are looking for the surface area of a cylinder, you need to find the area of two circles (the bases) and the area of the curved surface that makes up the side of the cylinder. The area of the curved surface is hard to visualize when it is rolled up. Picture a paper towel roll. It has a circular top and bottom. When you unroll a sheet of the paper towel, it is shaped like a rectangle. The area of the curved surface is the area of a rectangle with the same height as the cylinder, and the base measurement is the same as the circumference of the circle base.




$$\begin{aligned} \text{Surface area of a cylinder} &= \text{area of two circles} + \text{area of rectangle} \\ &= 2\pi r^2 + bh \\ &= 2\pi r^2 + 2\pi rh \end{aligned}$$


*Theorem:* The surface area (S.A.) of a cylinder is determined by finding the sum of the area of the bases and the product of the circumference times the height.

$$S.A. = 2\pi r^2 + 2\pi rh$$

**Examples:** Find the surface area of each cylinder. Use 3.14 for  $\pi$ .




$$\begin{aligned} S.A. &= 2\pi r^2 + 2\pi rh \\ S.A. &\approx 2(3.14)(4)^2 + 2(3.14)(4)(5) \\ S.A. &\approx 100.48 + 125.6 \\ S.A. &\approx 226.08 \text{ cm}^2 \end{aligned}$$



$$\begin{aligned} S.A. &= 2\pi r^2 + 2\pi rh \\ S.A. &\approx 2(3.14)(3)^2 + 2(3.14)(3)(10) \\ S.A. &\approx 56.52 + 188.4 \\ S.A. &\approx 244.92 \text{ ft.}^2 \end{aligned}$$


**Practice**

Find the surface area of each cylinder shown or described. Use 3.14 for  $\pi$ .



\_\_\_\_\_ **11.** Cylinder:  $r = 6$  ft.,  $h = 16$  ft.

\_\_\_\_\_ **12.** Cylinder:  $d = 18$  cm,  $h = 3$  cm




**► Volume of a Cylinder**

Similar to finding the volume of a prism, you can find the volume of a cylinder by finding the product of the area of the base and the height of the figure. Of course, the base of a cylinder is a circle, so you need to find the area of a circle times the height.

*Theorem:* The volume ( $V$ ) of a cylinder is the product of the area of the base ( $B$ ) and the height ( $h$ ).

$$V = Bh \text{ or } V = \pi r^2 h$$

**Examples:** Find the volume of each cylinder. Use 3.14 for  $\pi$ .




$$V = \pi r^2 h$$

$$V \approx (3.14)(3)^2(9)$$

$$V \approx (3.14)(9)(9)$$

$$V \approx 254.34 \text{ cm}^3$$



$$V = \pi r^2 h$$

$$V \approx (3.14)(6)^2(8)$$


$$V \approx (3.14)(36)(8)$$

$$V \approx 904.32 \text{ in.}^3$$

**Practice**

Find the volume of each cylinder shown or described. Use 3.14 for  $\pi$ .


\_\_\_\_\_ **13.**



\_\_\_\_\_ **15.** Cylinder:  $r = 5 \text{ in.}$ ,  $h = 24 \text{ in.}$


\_\_\_\_\_ **16.** Cylinder:  $d = 22 \text{ ft.}$ ,  $h = 12 \text{ ft.}$

\_\_\_\_\_ **14.**




**► Volume of a Cone**

A cone relates to a cylinder in the same way that a pyramid relates to a prism. If you have a cone and a cylinder with the same radius and height, then it would take three of the cones to fill the cylinder. In other words, the cone holds one-third the amount of the cylinder.




$$V = Bh \text{ or } V = \pi r^2 h$$



$$V = \frac{1}{3}Bh \text{ or } V = \frac{1}{3}\pi r^2 h$$

**Example:** Find the volume of the cone. Use 3.14 for  $\pi$ .



$$V = \frac{1}{3}\pi r^2 h$$


$$V \approx \frac{1}{3}(3.14)(3)^2(7)$$

$$V \approx 65.94 \text{ in.}^3$$

**Practice**

Find the volume of each cone shown or described. Use 3.14 for  $\pi$ .


\_\_\_\_\_ **17.**



\_\_\_\_\_ **19.** Cone:  $r = 3 \text{ cm}$ ,  $h = 9 \text{ cm}$

\_\_\_\_\_ **20.** Cone:  $d = 14 \text{ ft.}$ ,  $h = 5 \text{ ft.}$

\_\_\_\_\_ **18.**




► **Surface Area of a Sphere**

A *sphere* is the set of all points that are the same distance from some point called the center. A sphere is most likely to be called a ball. Try to find an old baseball and take the cover off of it. (Make sure it does not have Mark McGwire's autograph on it first!) When you lay out the cover of the ball, it roughly appears to be four circles. Recall that the formula for finding the area of a circle is  $A = \pi r^2$ .

*Theorem:* The surface area (S.A.) formula for a sphere is four times  $\pi$  times the radius squared.

$$S.A. = 4\pi r^2$$

**Example:** Find the surface area of the sphere. Use 3.14 for  $\pi$ .



$$S.A. = 4\pi r^2$$


$$S.A. \approx (4)(3.14)(2)^2$$

$$S.A. \approx 50.24 \text{ cm}^2$$


**Practice**

Find the surface area of each sphere shown or described. Use 3.14 for  $\pi$ .

\_\_\_\_\_ **21.**



\_\_\_\_\_ **22.**




\_\_\_\_\_ **23.** Sphere:  $r = 16 \text{ cm}$

\_\_\_\_\_ **24.** Sphere:  $r = 20 \text{ ft.}$

**► Volume of a Sphere**

If you were the filling balloons with helium, it would be important for you to know the volume of a sphere. To find the volume of a sphere, picture the sphere filled with numerous pyramids. The height of each pyramid represents the radius ( $r$ ) of the sphere. The sum of the areas of all the bases represents the surface area of the sphere.



$$\text{Volume of each pyramid} = \frac{1}{3}Bh$$

$$\text{Sum of the volumes} = n \times \frac{1}{3}Br$$

$$= \frac{1}{3}(nB)r$$

$$= \frac{1}{3}(4\pi r^2)r$$

$$= \frac{4}{3}\pi r^3$$


Substitute  $r$  for  $h$  of  $n$  pyramids

Substitute  $nB$  with the S.A. of a sphere

*Theorem:* The volume ( $V$ ) of a sphere is determined by the product of  $\frac{4}{3}\pi$  times the radius cubed.

$$V = \frac{4}{3}\pi r^3$$

**Example:** Find the volume of the sphere. Use 3.14 for  $\pi$ .



$$V = \frac{4}{3}\pi r^3$$


$$V \approx \frac{4}{3}(3.14)(9)^3$$

$$V \approx 3052.08 \text{ cm}^3$$

**Practice**

Find the volume of each sphere shown or described. Use 3.14 for  $\pi$ .


\_\_\_\_\_ **25.**



\_\_\_\_\_ **27.** Sphere:  $r = 12 \text{ m}$

\_\_\_\_\_ **28.** Sphere:  $d = 9 \text{ ft.}$

\_\_\_\_\_ **26.**



**Skill Building until Next Time**

Find an example of a cylinder—an oatmeal box would be a good choice. Measure the height and radius of the cylinder. Use the formulas in this lesson to find the surface area and volume of the cylinder.

LESSON

# 17




# Coordinate Geometry

## LESSON SUMMARY

In this lesson, you will learn to identify the  $x$ -axis,  $y$ -axis, the origin, and the four quadrants on a coordinate plane. You will also learn how to plot or graph points on a coordinate plane and name the coordinates of a point. The distance between two points will also be found using a formula.

If you have ever been the navigator on a road trip, then you have probably read a road map or grid map. A grid map uses a horizontal and vertical axis in a similar manner as a coordinate plane.

On a coordinate plane, the horizontal axis is called the  $x$ -axis. The vertical axis on the coordinate plane is called the  $y$ -axis. The point where the two axes cross is called the *origin*.






The two axes divide the coordinate plane into four regions, which are called quadrants. The quadrants are numbered counterclockwise beginning with the upper-right region. The coordinates  $(x,y)$  of a point are an ordered pair of numbers. The first number is the  $x$ -coordinate. The second number is the  $y$ -coordinate. The coordinates of the origin are  $(0,0)$ .

**Example:** Graph point  $A(-4,1)$  and point  $B(5,-3)$ . In which quadrant would you find each point?

**Solution:**




To graph point  $A(-4,1)$ , from the origin, go left 4 units and up 1. Label the point  $A$ . Point  $A$  is in quadrant II. To graph point  $B(5,-3)$ , start from the origin, go right 5 units and down 3. Label the point  $B$ . Point  $B$  is in quadrant IV.

Some points may be graphed on the axes also.

**Example:** Graph point  $C(2,0)$  and point  $D(0,-6)$ . On which axis will each point be?

**Solution:**



To graph point  $C$ , from the origin, go right 2 units, but do not move up or down. Label the point  $C$ . The point  $C$  is on the  $x$ -axis. To graph point  $D$ , from the origin, do not move right or left; move 6 units down. Label the point  $D$ . The point  $D$  is on the  $y$ -axis.

**Practice**

Draw a set of axes on graph paper. Graph and label each point.

**1.**  $A(-6,2)$

**7.**  $G(1,4)$

**2.**  $B(6,4)$

**8.**  $H(-2,5)$

**3.**  $C(4,-5)$

**9.**  $I(-1,-1)$

**4.**  $D(-5,-5)$

**10.**  $J(3,-2)$

**5.**  $E(0,6)$

**11.**  $K(2,0)$

**6.**  $F(-3,0)$

**12.**  $L(0,-5)$

Identify which quadrant or axis in which the point lies for the following points:

\_\_\_\_\_ **13.**  $A(-6,2)$

\_\_\_\_\_ **19.**  $G(1,4)$

\_\_\_\_\_ **14.**  $B(6,4)$

\_\_\_\_\_ **20.**  $H(-2,5)$

\_\_\_\_\_ **15.**  $C(4,-5)$

\_\_\_\_\_ **21.**  $I(-1,-1)$

\_\_\_\_\_ **16.**  $D(-5,-5)$

\_\_\_\_\_ **22.**  $J(3,-2)$

\_\_\_\_\_ **17.**  $E(0,6)$

\_\_\_\_\_ **23.**  $K(2,0)$


\_\_\_\_\_ **18.**  $F(-3,0)$

\_\_\_\_\_ **24.**  $L(0,-5)$

### ► Finding the Coordinates of a Point

Each point on the coordinate plane has its own unique ordered pair. You can think of an ordered pair as an address. Now that you have located a point, you can also find the coordinates of a point on a graph.


**Example:** State the coordinates of each point.



**Solution:**  $A(2,2)$   $B(-3,1)$   $C(0,4)$   $D(-4,-2)$   $E(2,-3)$   $F(5,0)$

### Practice

State the coordinates of each point.



\_\_\_\_\_ 25. *A*

\_\_\_\_\_ 26. *B*

\_\_\_\_\_ 27. *C*

\_\_\_\_\_ 28. *D*

\_\_\_\_\_ 29. *E*


\_\_\_\_\_ 30. *F*

\_\_\_\_\_ 31. *G*

\_\_\_\_\_ 32. *H*

### ► Finding the Distance between Two Points

You can easily count to find the distance between two points on a horizontal or vertical line. For example, in the following figure,  $\overline{XY} = 3$  and  $\overline{YZ} = 4$ . However, you cannot find  $\overline{XZ}$  simply by counting. Since  $\triangle XYZ$  is a right triangle, you can use the Pythagorean theorem.



$$XZ^2 = 3^2 + 4^2$$

$$XZ^2 = 9 + 16$$

$$XZ^2 = 25$$

$$XZ = 5$$

Of course, you won't be able to use the Pythagorean theorem all the time. However, you can use the following formula to find the distance between any two points.

*Theorem:* The distance  $d$  between any two points  $A(x_1, y_1)$  and  $B(x_2, y_2)$  is

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

**Example:** Find the distance between  $R(-3, 4)$  and  $S(-3, -7)$ .

**Solution:** It helps to label the coordinates of the points before you insert them into the formula.

$$\begin{array}{ll} x_1 y_1 & x_2 y_2 \\ R(-3, 4) & S(-3, -7) \end{array}$$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$d = \sqrt{(-3 - (-3))^2 + (-7 - 4)^2}$$

$$d = \sqrt{(-3 + 3)^2 + (-11)^2}$$

$$d = \sqrt{0^2 + 121}$$

$$d = \sqrt{121}$$

$$d = 11$$

**Practice**

Find the distance between the points with the given coordinates.

\_\_\_\_\_ **33.** (3,5) and (-1,5)

\_\_\_\_\_ **34.** (3,-2) and (-5,4)

\_\_\_\_\_ **35.** (-3,-8) and (-6,-4)

\_\_\_\_\_ **36.** (-1,-6) and (4,6)

**Skill Building until Next Time**

Examine a city map to see if there are any streets that cut diagonally across horizontal and vertical streets. If so, construct a coordinate plane for the map and determine the distance between two locations that are on the diagonal street.

LESSON

18



# The Slope of a Line


## LESSON SUMMARY

In this lesson, you will learn how to determine the slope of a line from its graph or from two points on the line. You will also learn how to tell by sight if the slope of a line is positive, negative, zero, or undefined.

**T**he *slope* of a line is the measure of its steepness. The slope of a line is determined by the ratio of its rise to run. When looking at a line on a coordinate grid, always count the run before you count the rise. When a line points up to the right, it has a positive slope. A line with a negative slope points up to the left.


**Examples:** Find the slope of each line.

(a)



**Solution:** Slope of  $\overline{VW} = \frac{\text{rise}}{\text{run}} = \frac{1}{3}$

(b)




**Solution:** Slope of  $\overline{XY} = \frac{\text{rise}}{\text{run}} = \frac{-3}{5}$


**Practice**

Find the slope of each line.


\_\_\_\_\_ 1.




\_\_\_\_\_ 4.




\_\_\_\_\_ 2.




\_\_\_\_\_ 5.



\_\_\_\_\_ 3.





\_\_\_\_\_ 6.







## ► Special Cases of Slope

You may be wondering what the slope is for horizontal and vertical lines—these two types of lines do not slant up toward the left or the right. They are special cases.




The horizontal line has a slope of zero, since the rise is equal to zero. (Remember that anything divided by zero equals zero, so  $\frac{\text{rise}}{\text{run}} = \frac{0}{x} = 0$ .) On the other hand, the vertical line has an undefined slope (sometimes said to have “no slope”). In this case, the run is equal to zero, so  $\frac{\text{rise}}{\text{run}} = \frac{y}{0}$ , and a number divided by zero is said to be *undefined*, since it doesn’t make sense to take something and divide it up into zero parts. You may think that zero slope and an undefined slope are the same thing, but ask yourself if you would prefer to drive on a road with a speed limit of zero or a road with no speed limit. You can see that these are very different. The following illustrations may help you remember which type of line—horizontal or vertical—has an undefined (no slope) or a slope of zero.



**Practice**

Tell whether the slope of each line is positive, negative, zero, or undefined (no slope).


7.



10.



8.



11.



9.



12.



**► Using a Formula**

You can also use a formula to determine the slope of a line containing two points, point  $A(x_1, y_1)$  and point  $B(x_2, y_2)$ . Here is the formula:

$$\text{slope} = \frac{y_2 - y_1}{x_2 - x_1}$$

**Example:** Find the slope of the line through  $P(-6, 5)$  and  $Q(-2, -1)$ .

**Solution:** Begin by labeling the coordinates before you insert the values into the formula.

$x_1 y_1$	$x_2 y_2$
$P(-6, 5)$	$Q(-2, -1)$

$$\text{Slope} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-1 - 5}{-2 - (-6)} = \frac{-6}{-2 + 6} = \frac{-6}{4} = \frac{-3}{2}$$

**Practice**

Use a formula to find the slope of the line through each pair of points.

\_\_\_\_\_ **13.** (0,0) and (5,6)

\_\_\_\_\_ **17.** (-2,2) and (-2,3)

\_\_\_\_\_ **14.** (-3,-2) and (-4,-3)

\_\_\_\_\_ **18.** (-3,0) and (-7,-3)

\_\_\_\_\_ **15.** (1,-3) and (-1,-3)


\_\_\_\_\_ **19.** (8,-5) and (-10,7)

\_\_\_\_\_ **16.** (5,-6) and (-2,8)

\_\_\_\_\_ **20.** (5,6) and (-1,6)

**Skill Building until Next Time**

Locate a set of steps near your home or school. Measure the rise and run of the set of steps. Use these measurements to find the slope of the steps.



LESSON

19



# The Equation of a Line

## LESSON SUMMARY

In this lesson, you will learn how to identify linear equations. You will also learn how to use the equation of a line to find points on the line. You will also determine if an ordered pair is on a line using the equation of the line.

**P**eople often make comparisons among numbers. For example, a nurse looks at a patient's temperature over a period of time. A salesman will compare commissions from sports cars and family vans. These types of comparisons can often be graphed in a straight line to make predictions about future events. This straight line may be created by using a linear equation.

### ► What Is a Linear Equation?

The standard form for a linear equation is  $Ax + By = C$ , where  $A$ ,  $B$ , and  $C$  are constants;  $A$  and  $B$  cannot be zero at the same time. Linear equations will not have exponents on the variables  $x$  and  $y$ . The product or quotient of variables is not found in linear equations. Take a look at the examples of linear and nonlinear equations on the next page.

Linear equations

$$2x + 3y = 7$$

$$x = -3$$

$$y = \frac{2}{3}x + 4$$

$$y = 7$$

Nonlinear equations

$$2x^2 + 3y = 7$$

$$\frac{-3}{y} = x$$

$$y^2 = \frac{2}{3}x + 4$$

$$xy = 7$$

### Practice

Identify each equation as linear or nonlinear.

1.  $x - 3y = 0$

2.  $3y^2 = 2x$

3.  $x^2 + y^2 = 49$

4.  $x + y = 15$

5.  $x + y = 15$

6.  $\frac{1}{y} = 2$

7.  $\frac{1}{3}x = 36$

8.  $4x - 7y = 28$

### ► Points on a Line

When you graph points on a coordinate plane, you can easily see if they could be connected to form a straight line. However, it is possible for you to determine if a point is on a line or satisfies the equation of a line without using a coordinate plane. If the ordered pair can replace  $x$  and  $y$ , and make a true statement, then the ordered pair is a point on the line.

**Examples:** Determine if the ordered pairs satisfy the linear equation  $2x - y = 4$ .

(a)  $(2,2)$

**Solution:**  $2x - y = 4$

$$2(2) - 2 \stackrel{?}{=} 4$$

$$4 - 2 \stackrel{?}{=} 4$$

$$2 \neq 4$$

no;  $(2,2)$  is not on the line  $2x - y = 4$ .

(b)  $(0,-4)$

**Solution:**  $2x - y = 4$

$$2(0) - (-4) \stackrel{?}{=} 4$$

$$0 + 4 \stackrel{?}{=} 4$$

$$4 = 4$$

yes;  $(0,-4)$  is on the line  $2x - y = 4$ .

**Practice**

Determine if the ordered pairs satisfy the linear equation  $2x + 5y = 10$ .

- 9. (0,-2) 13. (15,4)
- 10. (10,1) 14. (3,-1)
- 11.  $(\frac{5}{2}, 3)$  15.  $(-\frac{5}{2}, -3)$
- 12. (-5,-4) 16. (5,0)

► **Graphing Linear Equations**

You can graph any linear equation you want by choosing several  $x$  or  $y$  coordinates to substitute into the original equation, and then solve the equation to find the other variable. This is easier if you organize your work into a table. You could choose any values you want and get a solution that would be an ordered pair or point on the line, but for this example, let's use the  $x$ -values.


**Example:** Find three ordered pairs that satisfy the equation. Graph each equation.

$y = x + 4$

**Solution:**

$x$	$x + 4$	$y$	$(x, y)$
-2			
0			
2			

$x$	$x + 4$	$y$	$(x, y)$
-2	$-2 + 4$	2	$(-2, 2)$
0	$0 + 4$	4	$(0, 4)$
2	$2 + 4$	6	$(2, 6)$



**Practice**

Find three ordered pairs that satisfy each equation. Then graph each equation.

17.  $y = 2x + 3$

$x$	$2x + 3$	$y$	$(x, y)$
2			
0			
-2			

18.  $y = -x$

$x$	$-x$	$y$	$(x, y)$
2			
0			
-2			

19.  $y = 2x + 4$

$x$	$-2x + 4$	$y$	$(x, y)$
1			
0			
-1			

20.  $y = 4x - 5$

$x$	$4x - 5$	$y$	$(x, y)$
1			
0			
-1			

**Skill Building until Next Time**

The distance that a lightning flash is from you and the time in seconds that it takes for you to hear the thunder clap can be written in the following linear equation:  $d = \frac{t}{5}$ . Find at least three ordered pairs  $(t, d)$  that satisfy the formula and graph the line.

LESSON

20



# Trigonometry Basics


## LESSON SUMMARY

In this lesson, you will learn how to write the trigonometric ratios sine, cosine, and tangent for a right triangle. You will learn how to use the trigonometry ratios to find unknown angle or side measurements.

**P**roperties of similar right triangles are the basis for trigonometry. Measurements sometimes cannot be made by using a measuring tape. For example, the distance from a ship to an airplane can't be measured this way; however, it could be found by using trigonometric ratios. The word *trigonometry* comes from the Greek language; it means "triangle measurement."

Recall that the hypotenuse is the side of the triangle across from the right angle. The other two sides of the triangle are called legs. These two legs have special names in relation to a given angle: adjacent leg and opposite leg. The word *adjacent* means *beside*, so the adjacent leg is the leg beside the angle. The opposite leg is across from the angle. You can see in the figure on the next page how the legs are named depending on which acute angle is selected.






Sine, cosine, and tangent ratios for the acute angle  $A$  in a right triangle are as follows:

$$\sin A = \frac{\text{opposite leg}}{\text{hypotenuse}}$$

$$\cos A = \frac{\text{adjacent leg}}{\text{hypotenuse}}$$

$$\tan A = \frac{\text{opposite leg}}{\text{adjacent leg}}$$

**Examples:** Express each ratio as a decimal to the nearest thousandth.



- (a)  $\sin A$ ,  $\cos A$ ,  $\tan A$   
 (b)  $\sin B$ ,  $\cos B$ ,  $\tan B$

**Solution:**

$$(a) \quad \sin A = \frac{\text{opp}}{\text{hyp}} = \frac{4}{5} = 0.800$$

$$\cos A = \frac{\text{adj}}{\text{hyp}} = \frac{3}{5} = 0.600$$

$$\tan A = \frac{\text{opp}}{\text{adj}} = \frac{4}{3} \approx 1.333$$


$$(b) \quad \sin B = \frac{\text{opp}}{\text{hyp}} = \frac{3}{5} = 0.600$$

$$\cos B = \frac{\text{adj}}{\text{hyp}} = \frac{4}{5} = 0.800$$

$$\tan B = \frac{\text{opp}}{\text{adj}} = \frac{3}{4} = 0.750$$


**Practice**

Express each ratio as a decimal to the nearest thousandth. Use the following figure to answer practice problems 1–6.



- \_\_\_\_\_ **1.**  $\sin Y$
- \_\_\_\_\_ **2.**  $\cos Y$
- \_\_\_\_\_ **3.**  $\tan Y$
- \_\_\_\_\_ **4.**  $\sin Z$
- \_\_\_\_\_ **5.**  $\cos Z$
- \_\_\_\_\_ **6.**  $\tan Z$

Use the following figure to answer practice problems 7–12.



- \_\_\_\_\_ **7.**  $\sin L$
- \_\_\_\_\_ **8.**  $\cos L$
- \_\_\_\_\_ **9.**  $\tan L$
- \_\_\_\_\_ **10.**  $\sin N$
- \_\_\_\_\_ **11.**  $\cos N$
- \_\_\_\_\_ **12.**  $\tan N$

## ► Using a Trigonometric Table

Trigonometric ratios for all acute angles are commonly listed in tables. Scientific calculators also have functions for the trigonometric ratios. Consult your calculator handbook to make sure you have your calculator in the *degree*, and not the *radian* setting. Part of a trigonometric table is given here.

ANGLE	SIN	COS	TAN
16°	0.276	0.961	0.287
17°	0.292	0.956	0.306
18°	0.309	0.951	0.325
19°	0.326	0.946	0.344
20°	0.342	0.940	0.364
21°	0.358	0.934	0.384
22°	0.375	0.927	0.404
23°	0.391	0.921	0.424
24°	0.407	0.914	0.445
25°	0.423	0.906	0.466
26°	0.438	0.899	0.488
27°	0.454	0.891	0.510
28°	0.470	0.883	0.532
29°	0.485	0.875	0.554
30°	0.500	0.866	0.577
31°	0.515	0.857	0.601
32°	0.530	0.848	0.625
33°	0.545	0.839	0.649
34°	0.559	0.829	0.675
35°	0.574	0.819	0.700
36°	0.588	0.809	0.727
37°	0.602	0.799	0.754
38°	0.616	0.788	0.781
39°	0.629	0.777	0.810
40°	0.643	0.766	0.839
41°	0.656	0.755	0.869
42°	0.669	0.743	0.900
43°	0.682	0.731	0.933
44°	0.695	0.719	0.966
45°	0.707	0.707	1.000

**Example:** Find each value.

(a)  $\cos 44^\circ$

(b)  $\tan 42^\circ$

**Solution:**

(a)  $\cos 44^\circ = 0.719$

(b)  $\tan 42^\circ = 0.900$

**Example:** Find  $m\angle A$ .

(a)  $\sin A = 0.656$

(b)  $\cos A = 0.731$

**Solution:**

(a)  $m\angle A = 41^\circ$

(b)  $m\angle A = 43^\circ$

**Practice**

Use the trigonometric table or a scientific calculator to find each value to the nearest thousandth or each angle measurement rounded to the nearest degree.

\_\_\_\_\_ **13.**  $\sin 18^\circ$

\_\_\_\_\_ **19.**  $\sin A = 0.485$

\_\_\_\_\_ **14.**  $\sin 32^\circ$

\_\_\_\_\_ **20.**  $\cos A = 0.743$

\_\_\_\_\_ **15.**  $\cos 27^\circ$

\_\_\_\_\_ **21.**  $\tan A = 0.384$

\_\_\_\_\_ **16.**  $\cos 40^\circ$

\_\_\_\_\_ **22.**  $\cos A = 0.788$

\_\_\_\_\_ **17.**  $\tan 36^\circ$

\_\_\_\_\_ **23.**  $\sin A = 0.375$


\_\_\_\_\_ **18.**  $\tan 17^\circ$

\_\_\_\_\_ **24.**  $\tan A = 0.306$

**► Finding the Measure of an Acute Angle**

The trigonometric ratio used to find the measure of an acute angle of a right triangle depends on which side lengths are known.

**Example:** Find  $m\angle A$ .



**Solution:** The  $\sin A$  involves the two lengths known.

$$\sin A = \frac{\text{opp}}{\text{hyp}} = \frac{5}{16} \approx 0.313$$

Note that the  $\approx$  symbol is used in the previous solution because the decimal 0.313 has been rounded to the nearest thousandths place.

Using the sin column of a trigonometric table, you'll find:

$$\sin 18^\circ = 0.309 \text{ } \} \text{ difference } 0.004$$


$$\sin A \approx 0.313$$

$$\sin 19^\circ = 0.326 \text{ } \} \text{ difference } 0.013$$

Since 0.313 is closer to 0.309,  $m\angle A \approx 18^\circ$  to the nearest degree.

A scientific calculator typically includes a button that reads  $\sin^{-1}$ , which means *inverse sine*, and can be used to find an angle measure when the sine is known.

**Example:** Find  $m\angle C$ .



**Solution:** The tan C involves the two lengths known.

$$\tan C = \frac{\text{opp}}{\text{adj}} = \frac{8}{11} \approx 0.727$$

Using the tan column, you'll find:

$$\tan 36 = 0.727$$


Therefore,  $m\angle C = 36^\circ$ .

Again, the  $\tan^{-1}$  button on a scientific calculator can be used to find an angle measure when its tangent is known.


**Practice**

Find the measure of  $\angle A$  to the nearest degree.


\_\_\_\_\_ 25.




\_\_\_\_\_ 27.



\_\_\_\_\_ 26.



\_\_\_\_\_ 28.




## ► Finding the Measure of a Side

The trigonometric ratio used to find the length of a side of a right triangle depends on which side length and angle are known.


**Example:** Find the value of  $x$  to the nearest tenth.

(a)



**Solution:**  $\sin 35 = \frac{x}{10}$   
 $0.574 = \frac{x}{10}$   
 $x = 10(0.574)$   
 $x \approx 5.7$





(b)



**Solution:**  $\cos 28 = \frac{9}{x}$   
 $0.883 = \frac{9}{x}$   
 $x = \frac{9}{0.883}$   
 $x \approx 10.2$


**Practice**

Find the value of  $x$  to the nearest tenth.



**Skill Building until Next Time**

Find a set of steps near your home or school. Measure the length of the hypotenuse and the adjacent leg. Then use cosine to find the angle of elevation of the steps. There are building codes that limit the angle of elevation of a set of steps to help prevent accidents.



# Posttest



**N**ow that you have completed all the lessons, it is time to show off your new skills. Take the posttest in this chapter to see how much your geometry skills have improved. The posttest has 50 multiple-choice questions covering the topics you studied in this book. While the format of the posttest is similar to that of the pretest, the questions are different.

After you complete the posttest, check your answers with the answer key at the end of the book. Along with each answer is a number that tells you which lesson of this book teaches you about the geometry skills needed for that question. If you still have weak areas, go back and rework the applicable lessons.





LEARNINGEXPRESS ANSWER SHEET

- 1. (a) (b) (c) (d)
- 2. (a) (b) (c) (d)
- 3. (a) (b) (c) (d)
- 4. (a) (b) (c) (d)
- 5. (a) (b) (c) (d)
- 6. (a) (b) (c) (d)
- 7. (a) (b) (c) (d)
- 8. (a) (b) (c) (d)
- 9. (a) (b) (c) (d)
- 10. (a) (b) (c) (d)
- 11. (a) (b) (c) (d)
- 12. (a) (b) (c) (d)
- 13. (a) (b) (c) (d)
- 14. (a) (b) (c) (d)
- 15. (a) (b) (c) (d)
- 16. (a) (b) (c) (d)
- 17. (a) (b) (c) (d)


- 18. (a) (b) (c) (d)
- 19. (a) (b) (c) (d)
- 20. (a) (b) (c) (d)
- 21. (a) (b) (c) (d)
- 22. (a) (b) (c) (d)
- 23. (a) (b) (c) (d)
- 24. (a) (b) (c) (d)
- 25. (a) (b) (c) (d)
- 26. (a) (b) (c) (d)
- 27. (a) (b) (c) (d)
- 28. (a) (b) (c) (d)
- 29. (a) (b) (c) (d)
- 30. (a) (b) (c) (d)
- 31. (a) (b) (c) (d)
- 32. (a) (b) (c) (d)
- 33. (a) (b) (c) (d)
- 34. (a) (b) (c) (d)

- 35. (a) (b) (c) (d)
- 36. (a) (b) (c) (d)
- 37. (a) (b) (c) (d)
- 38. (a) (b) (c) (d)
- 39. (a) (b) (c) (d)
- 40. (a) (b) (c) (d)
- 41. (a) (b) (c) (d)
- 42. (a) (b) (c) (d)
- 43. (a) (b) (c) (d)
- 44. (a) (b) (c) (d)
- 45. (a) (b) (c) (d)
- 46. (a) (b) (c) (d)
- 47. (a) (b) (c) (d)
- 48. (a) (b) (c) (d)
- 49. (a) (b) (c) (d)
- 50. (a) (b) (c) (d)



► Posttest

1. Which pair of angles are supplementary?




- a.  $\angle 4$  and  $\angle 5$
- b.  $\angle 4$  and  $\angle 8$
- c.  $\angle 4$  and  $\angle 6$
- d.  $\angle 4$  and  $\angle 1$

2. What is the range of degrees for an acute angle?


- a. greater than 0, less than 90
- b. greater than 90, less than 180
- c. greater than 0, less than 180
- d. greater than 90, less than 360

3. In the figure,  $\angle XWZ = 130^\circ$  and  $\angle YWZ = 70^\circ$ .  
What is the measure of  $\angle XWY$ ?



- a.  $60^\circ$
- b.  $55^\circ$
- c.  $50^\circ$
- d.  $65^\circ$

4. What is the measure of  $\angle MOL$ ?




- a.  $130^\circ$
- b.  $50^\circ$
- c.  $40^\circ$
- d.  $120^\circ$

5. Classify  $\triangle MNO$  with  $\angle M = 25^\circ$ ,  $\angle N = 65^\circ$ , and  $\angle O = 90^\circ$ .

- a. acute
- b. right
- c. obtuse
- d. equiangular

6. Which postulate would you use to prove  $\triangle MNO \cong \triangle RST$ ?




- a. AAA
- b. SAS
- c. ASA
- d. SSS

7. Three sides of a triangle are 6, 8, and 10; what type of triangle is it?

- a. acute
- b. right
- c. obtuse
- d. straight

8. Which of the following is not an acceptable name for the pentagon?




- a. MESHO
  - b. SEMOH
  - c. HEMOS
  - d. HOMES
9. Which of the following is a property of rectangles?
- a. The diagonals are congruent.
  - b. Opposite sides are congruent.
  - c. Opposite sides are parallel.
  - d. all of the above
10. Solve for  $x$ :  $\frac{11}{25} = \frac{x}{100}$ .
- a. 44
  - b. 4
  - c. 11
  - d. 25
11. Find the perimeter of a rectangle with base 5 cm and height 20 cm.
- a. 100 cm
  - b. 25 cm
  - c. 50 cm
  - d. 55 cm
12. Find the area of a triangle with a base that measures 35 feet and height 10 feet.
- a.  $350 \text{ ft.}^2$
  - b.  $70 \text{ ft.}^2$
  - c.  $3.5 \text{ ft.}^2$
  - d.  $175 \text{ ft.}^2$

13. Find the surface area of a cube that measures 5 inches on an edge. Use  $S.A. = 6e^2$ .

- a.  $300 \text{ in.}^2$
- b.  $900 \text{ in.}^2$
- c.  $150 \text{ in.}^2$
- d.  $30 \text{ in.}^2$

14. Find the volume of the pyramid.




- a.  $60 \text{ m}^3$
  - b.  $180 \text{ m}^3$
  - c.  $30 \text{ m}^3$
  - d.  $36 \text{ m}^3$
15. Find the volume of a cone with radius 9 cm and height 2 cm. Use 3.14 for  $\pi$ .
- a.  $508.68 \text{ cm}^3$
  - b.  $3194.51 \text{ cm}^3$
  - c.  $56.52 \text{ cm}^3$
  - d.  $169.56 \text{ cm}^3$
16. In which quadrant will you graph the point  $(-4,8)$ ?
- a. I
  - b. II
  - c. III
  - d. IV
17. Find the slope of the line that passes through the points  $(1,2)$  and  $(3,3)$ .
- a.  $\frac{4}{5}$
  - b. 2
  - c.  $\frac{1}{2}$
  - d.  $\frac{5}{4}$

18. Which ordered pair does not satisfy the equation

$$2x - y = 6?$$


- a. (3,0)
- b. (0,-6)
- c. (1,-4)
- d. (2,6)

19. What is the  $\sin B$  for this figure?




- a.  $\frac{4}{3}$
- b.  $\frac{3}{4}$
- c.  $\frac{3}{5}$
- d.  $\frac{4}{5}$

20. Which of the following sets of points are collinear?




- a. A, B, C
- b. B, C, D
- c. B, C, E
- d. D, C, A

21. Which pairs of angles are congruent?



- a.  $\angle 1$  and  $\angle 2$
- b.  $\angle 1$  and  $\angle 4$
- c.  $\angle 1$  and  $\angle 3$
- d.  $\angle 1$  and  $\angle 7$

22. What is the correct classification for the angle?




- a. acute
- b. right
- c. obtuse
- d. straight

23. What type of angle is shaped like a corner of a piece of paper?


- a. acute
- b. right
- c. obtuse
- d. straight

24. What is the measurement of  $\angle AOC$ ?




- a.  $20^\circ$
- b.  $80^\circ$
- c.  $60^\circ$
- d.  $85^\circ$

25. In the figure  $\angle OMN = 45^\circ$ , what is the measurement of  $\angle LMO$ ?




- a.  $120^\circ$
- b.  $125^\circ$
- c.  $130^\circ$
- d.  $135^\circ$

26. What type of angles are  $\angle MON$  and  $\angle LOP$ ?




- a. vertical
- b. complementary
- c. supplementary
- d. adjacent

27. Classify the triangle in the figure by its angles.




- a. acute
- b. right
- c. obtuse
- d. equiangular

28.  $\triangle DRY$  is an isosceles triangle. What is the best name for  $\angle D$ ?




- a. leg
- b. vertex angle
- c. base angle
- d. base

29. Which postulate would you use to prove  $\triangle EFH \cong \triangle GFH$ ?



- a. AAA
- b. SAS
- c. ASA
- d. SSS

30. Find the missing length.



- a. 2
- b. 14
- c. 12
- d. 10

31. Three sides of a triangle are 7, 7, and 10; what type of triangle is it?


- a. acute
- b. right
- c. obtuse
- d. straight

32. What is the sum of the measures of the interior angles of a convex quadrilateral?

- a.  $45^\circ$
- b.  $90^\circ$
- c.  $180^\circ$
- d.  $360^\circ$


- 33.** Which of the following is not a property of parallelograms?
- Opposite sides of a parallelogram are congruent.
  - Opposite sides of a parallelogram are parallel.
  - Opposite angles of a parallelogram are congruent.
  - Diagonals of a parallelogram are congruent.

- 34.**  $ABCD$  is a square. Which of the following is not true?



- $m\angle BCD = 90^\circ$
  - $\overline{AB} \cong \overline{AC}$
  - $m\angle AXB = 90^\circ$
  - $\overline{AC} \cong \overline{BD}$
- 35.** Solve for  $y$ :  $\frac{12}{y} = \frac{36}{3}$ .
- 12
  - 3
  - 1
  - 36
- 36.** Find the perimeter of a regular pentagon that measures 3 ft. on each side.
- 3 ft.
  - 9 ft.
  - 12 ft.
  - 15 ft.
- 37.** Find the length of a side of an equilateral triangle whose perimeter is 42 m.
- 126 m
  - 21 m
  - 14 m
  - 7 m

- 38.** Find the area of the trapezoid in the figure.



- $22 \text{ cm}^2$
  - $72 \text{ cm}^2$
  - $36 \text{ cm}^2$
  - $18 \text{ cm}^2$
- 39.** Find the surface area of a right rectangular prism with length 4 m, width 2 m, and height 8 m. Use the formula  $S.A. = 2(lw + wh + lh)$ .
- $64 \text{ m}^2$
  - $112 \text{ m}^2$
  - $56 \text{ m}^2$
  - $32 \text{ m}^2$
- 40.** How many faces does a cube have?
- 4
  - 6
  - 8
  - 12
- 41.** Find the volume of a pyramid whose base has an area of  $17 \text{ in.}^2$  and whose height is 3 in.
- $51 \text{ in.}^3$
  - $17 \text{ in.}^3$
  - $3 \text{ in.}^3$
  - $20 \text{ in.}^3$
- 42.** Find the volume of a cylinder with radius 5 cm and height 3 cm. Use 3.14 for  $\pi$ .
- $47.1 \text{ cm}^3$
  - $78.5 \text{ cm}^3$
  - $141.3 \text{ cm}^3$
  - $235.5 \text{ cm}^3$




43. Find the surface area of a sphere with an 8-inch radius. Use 3.14 for  $\pi$ .

- a. 25.12 in.<sup>2</sup>
- b. 631.01 in.<sup>2</sup>
- c. 803.84 in.<sup>2</sup>
- d. 200.96 in.<sup>2</sup>

44. Find the distance between (1,2) and (3,2). Use  $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$ .

- a. 1
- b. 2
- c. 3
- d. 4

45. Find the slope of the line in the figure.



- a.  $\frac{2}{3}$
- b.  $\frac{3}{2}$
- c.  $-\frac{2}{3}$
- d.  $-\frac{3}{2}$

46. Find the slope of the line that passes through the points (0,0) and (-1,-4).

- a. -4
- b. 4
- c.  $\frac{1}{4}$
- d.  $-\frac{1}{4}$


47. If the  $x$ -value is 4, the  $y$ -value is \_\_\_\_ in the equation  $-x + y = 5$ .

- a. 9
- b. 1
- c. 5
- d. -1

48. What is the trigonometric ratio for sine?


- a.  $\frac{\text{opposite leg}}{\text{hypotenuse}}$
- b.  $\frac{\text{adjacent leg}}{\text{hypotenuse}}$
- c.  $\frac{\text{opposite leg}}{\text{adjacent leg}}$
- d.  $\frac{\text{adjacent leg}}{\text{opposite leg}}$

49. In the following figure, what is the  $\tan A$ ?



- a.  $\frac{4}{3}$
- b.  $\frac{3}{4}$
- c.  $\frac{3}{5}$
- d.  $\frac{4}{5}$

50. Which of the following sets contains noncollinear points?



- a. A, C
- b. C, D
- c. A, C, D
- d. A, B, D

# Answer Key





## ► Pretest

If you miss any of the answers, you can find help for that kind of question in the lesson shown to the right of the answer.

- |                       |                        |
|-----------------------|------------------------|
| <b>1. d</b> Lesson 1  | <b>21. d</b> Lesson 9  |
| <b>2. a</b> Lesson 1  | <b>22. a</b> Lesson 9  |
| <b>3. c</b> Lesson 1  | <b>23. b</b> Lesson 9  |
| <b>4. c</b> Lesson 2  | <b>24. a</b> Lesson 10 |
| <b>5. d</b> Lesson 2  | <b>25. c</b> Lesson 10 |
| <b>6. d</b> Lesson 3  | <b>26. c</b> Lesson 11 |
| <b>7. b</b> Lesson 3  | <b>27. b</b> Lesson 11 |
| <b>8. a</b> Lesson 3  | <b>28. d</b> Lesson 11 |
| <b>9. d</b> Lesson 4  | <b>29. b</b> Lesson 12 |
| <b>10. c</b> Lesson 4 | <b>30. c</b> Lesson 12 |
| <b>11. b</b> Lesson 5 | <b>31. a</b> Lesson 13 |
| <b>12. b</b> Lesson 5 | <b>32. a</b> Lesson 13 |
| <b>13. a</b> Lesson 5 | <b>33. b</b> Lesson 13 |
| <b>14. a</b> Lesson 6 | <b>34. d</b> Lesson 14 |
| <b>15. b</b> Lesson 6 | <b>35. a</b> Lesson 14 |
| <b>16. c</b> Lesson 7 | <b>36. c</b> Lesson 15 |
| <b>17. a</b> Lesson 7 | <b>37. c</b> Lesson 15 |
| <b>18. d</b> Lesson 7 | <b>38. d</b> Lesson 15 |
| <b>19. c</b> Lesson 8 | <b>39. d</b> Lesson 16 |
| <b>20. c</b> Lesson 8 | <b>40. c</b> Lesson 16 |

41. d Lesson 17  
 42. b Lesson 17  
 43. a Lesson 17  
 44. a Lesson 18  
 45. d Lesson 18  
 46. d Lesson 19  
 47. d Lesson 19  
 48. d Lesson 19  
 49. c Lesson 20  
 50. b Lesson 20

### ► Lesson 1

- Yes; there are countless points on any line.
- There are many examples of a point in everyday life. Here are a few: pupil of the eye, the dot above the letter “i,” the location of a city on a map, and a freckle.
- Lines, segments, rays, and planes are made up of a series of points.
- $\overleftrightarrow{XY}$ ,  $\overleftrightarrow{YZ}$ ,  $\overleftrightarrow{XZ}$ ,  $\overleftrightarrow{YX}$ ,  $\overleftrightarrow{ZY}$ ,  $\overleftrightarrow{ZX}$
- An infinite number of points are on a line.
- The notation for a line has two arrowheads because a line extends forever in both directions.
- $\overrightarrow{SR}$ ;  $\overrightarrow{ST}$
- The endpoint is the beginning of a ray.
- They are different because they have different endpoints and extend in different directions.
- $\overline{LM}$ ;  $\overline{MN}$ ;  $\overline{NP}$ ;  $\overline{LN}$ ;  $\overline{MP}$ ;  $\overline{LP}$
- Line segments do not extend indefinitely. They have starting points and stopping points.
- An infinite number of points are on a line segment.
- A line has no endpoints; a ray has one endpoint.
- A line and a ray both have an infinite set of points.
- A ray extends indefinitely in one direction, but a segment has two endpoints.
- Both have an infinite set of points.
- A plane has two dimensions; a line has one dimension.
- No, any two points can be connected to form a line.
- Yes, a third point could be off the line.
- Yes, coplanar points can be noncollinear because two points could be on one line with a third point being in the same plane but not on the same line.
- Yes, collinear points must be coplanar because if a line is in a plane, then all points on that line are in the same plane.
- yes
- no
- yes
- Yes; remember that any two points determine a line, even if it is not drawn.
- yes
- no
- no
- Yes; remember that any three noncollinear points determine a plane, even if it is not drawn.
- true
- false
- true
- False; sometimes they do, but not always.
- 
- Here is one possible answer:  

- Not possible; any three points are coplanar.

### ► Lesson 2

- Vertex:  $Y$ ; sides:  $\overrightarrow{YX}$ ;  $\overrightarrow{YZ}$
- Vertex:  $J$ ; sides:  $\overrightarrow{JA}$ ;  $\overrightarrow{JT}$
- $\angle KBT$ ,  $\angle TBK$ ,  $\angle B$ ,  $\angle I$
- $\angle PON$  ( $\angle NOP$ ) and  $\angle POM$  ( $\angle MOP$ )
- $\angle MON$  ( $\angle NOM$ )
- $\overrightarrow{ON}$  and  $\overrightarrow{OM}$
- More than one angle uses the letter  $O$  as a vertex; therefore, no one would know which  $\angle O$  you meant.

8. No; they don't share the same endpoint.
9. No; they may not form a line.
10.  $\angle KOL$ ;  $\angle JOK$ ;  $\angle NOM$
11.  $\angle NOK$ ;  $\angle MOJ$ ;  $\angle MOL$
12.  $\angle NOL$ ;  $\angle MOK$
13.  $63^\circ$  ( $180^\circ - 27^\circ - 90^\circ$ )
14. right
15. straight
16. acute
17. obtuse
18. acute
19. straight
20. obtuse
21. right
22. acute
23. obtuse
24. right
25. straight


### ► Lesson 3

1. Yes; line  $d$  cuts across lines  $t$  and  $r$ .
2. Yes; line  $y$  cuts across lines  $t$  and  $r$ .
3. No; line  $t$  intersects lines  $d$  and  $y$  at the same point, not two different points.
4. Yes; line  $r$  cuts across lines  $d$  and  $y$  at two different points.
5. true
6. False; the symbol means perpendicular.
7. False; transversals can be perpendicular, but they do not have to be perpendicular.
8. always
9. never
10. sometimes
11. never
12. never
13. Corresponding angles; congruent
14. Same-side interior angles; supplementary
15. Corresponding angles; congruent
16. Alternate interior angles; congruent


17.  $100^\circ$
18.  $109^\circ$
19.  $110^\circ$
20.  $55^\circ$
21. They are both pairs of congruent angles when formed by parallel lines and a transversal.
22. Alternate interior angles are both "inside" the parallel lines. Corresponding angles are a pair of angles with one angle "inside" the parallel lines and one "outside" the parallel lines.
23. They are both pairs of interior angles.
24. Same-side interior angles are supplementary. Alternate interior angles are congruent.
25. Both pairs of angles are on the same side of the transversal.
26. Same-side interior angles are supplementary. Corresponding angles are congruent. Also, both same-side interior angles are interior while corresponding angles have one angle "inside" and one "outside" the parallel lines.

### ► Lesson 4


1.  $20^\circ$
2.  $75^\circ$
3.  $100^\circ$
4.  $135^\circ$
5.  $55^\circ$
6.  $80^\circ$
7.  $115^\circ$
8.  $160^\circ$
9.  $25^\circ$
10.  $60^\circ$
- 11.




12.




13.



14.



15.

16.  $100^\circ$ 17.  $50^\circ$ 18.  $138^\circ$ 19.  $35^\circ$ 20.  $55^\circ$ 21.  $90^\circ$ 22.  $180^\circ$ 23.  $125^\circ$ 24.  $145^\circ$ 25.  $35^\circ$ 

26. 1,600 mils

27. 2,400 mils

28. 800 mils

29. 3,200 mils

## ► Lesson 5

1.  $46^\circ$ 2.  $22^\circ$ 3.  $10^\circ$ 4.  $65^\circ$ 5.  $87^\circ$ 6.  $102^\circ$ 7.  $50^\circ$ 8.  $120^\circ$ 9.  $25^\circ$ 10.  $179^\circ$ 11.  $\angle AOE$  and  $\angle COD$ 12.  $\angle BOC$  and  $\angle COD$ 13.  $\angle AOE$  and  $\angle COD$ ;  $\angle EOD$  and  $\angle AOC$ 

14. true

15. false

16. false

17. true

18. true

19. true

20. false

21. true

22. true

23. false

24. false

25. false

26.  $\angle 2$  and  $\angle 5$ ;  $\angle 3$  and  $\angle 5$ ;  $\angle 1$  and  $\angle 4$ 

## ► Lesson 6

1. equilateral

2. isosceles

3. scalene

4. scalene

5. isosceles

6. scalene

7. equilateral

8. legs:  $\overline{XY}$  and  $\overline{YZ}$ vertex angle:  $\angle Y$

base angles:  $\angle X$  and  $\angle Z$

base:  $\overline{XZ}$

9. legs:  $\overline{RS}$  and  $\overline{ST}$

vertex angle:  $\angle S$

base angles:  $\angle R$  and  $\angle T$

base:  $\overline{RT}$

10. legs:  $\overline{DE}$  and  $\overline{EF}$

vertex angle:  $\angle E$

base angles:  $\angle D$  and  $\angle F$

base:  $\overline{DF}$

11. acute

12. equiangular and acute

13. right

14. obtuse

15. obtuse

16. acute

17. right

18. true

19. true

20. false

21. false

22. true

23. true

24. false

25. true

### ► Lesson 7

1.  $\angle R$

2.  $\angle J$

3.  $\angle K$

4.  $\overline{PQ}$

5.  $\overline{MK}$

6.  $\overline{JM}$

7.  $\triangle FJH$

8.  $\triangle HGF$

9.  $\triangle FHG$

10.  $\triangle GFH$

11.  $\overline{RS}$

12.  $\overline{VS}$

13.  $\overline{RT}$

14. yes

15. no

16. vertical angles

17. yes

18.  $\overline{AC}$

19.  $\overline{CD}$

20. no (should be  $\triangle ACB \cong \triangle ECD$ )

21.  $\angle QRS$

22.  $\angle TSR$

23. yes

24. no

25. yes

26. SAS

27. ASA

28. ASA

29. SSS

30. SAS

### ► Lesson 8

1.  $\overline{AB}$  and  $\overline{BC}$

2.  $\overline{AC}$

3.  $\overline{PR}$  and  $\overline{RQ}$

4.  $\overline{PQ}$

5.  $\overline{HL}$  and  $\overline{LM}$

6.  $\overline{HM}$

7.

Number	Square	Number	Square
1	1	9	81
2	4	10	100
3	9	11	121
4	16	12	144
5	25	13	169
6	36	14	196
7	49	15	225
8	64	16	256

8. 17

9. 9

10. 8

- 11.** 5  
**12.** 40  
**13.** 25  
**14.** yes  
**15.** no  
**16.** yes  
**17.** no  
**18.** yes  
**19.** yes  
**20.** right  
**21.** acute  
**22.** obtuse  
**23.** obtuse  
**24.** right  
**25.** acute  
**26.** obtuse  
**27.** When the ladder is placed 5 feet from the building, the ladder extends a little over 17 feet up the building. When the ladder is placed 3 feet from the building, it reaches about  $17\frac{3}{4}$  feet up the building. It would be impractical to place the ladder that close or even closer because it would not be stable. You would not go very far up the ladder before you would be falling back down!

## ► Lesson 9

- 1.** not a polygon  
**2.** convex polygon  
**3.** concave polygon  
**4.** convex polygon  
**5.** not a polygon  
**6.** convex polygon  
**7.** concave polygon  
**8.** not a polygon  
**9.** not a polygon  
**10.** quadrilateral  
**11.** triangle  
**12.** pentagon  
**13.** heptagon  
**14.** hexagon  
**15.** octagon

- 16.** yes  
**17.** no  
**18.** yes  
**19.** yes  
**20.** no  
**21.** no  
**22.**

<i>Number of sides</i>	6	10	14	16
<i>Interior ∠ sum</i>	720	1,440	2,160	2,520
<i>Exterior ∠ sum</i>	360	360	360	360

- 23.** Home plate is a pentagon, so use the formula with  $n = 5$ .  
 $S = 180(n - 2)$   
 $S = 180(5 - 2)$   
 $S = 180(3)$   
 $S = 540$   
 $3(\text{right angles}) + 2(\text{obtuse angles}) = 540$   
 $3(90) + 2(\text{obtuse angles}) = 540$   
 $270 + 2(\text{obtuse angles}) = 540$   
 $-270$  (subtract the 3 right  $\angle$ 's to see how much of the original 540 is left)  
 $2(\text{obtuse angles}) = 270$   
 obtuse angle = 135 (divide by 2 because the two angles share the 270 evenly)  
 Therefore, each obtuse angle equals  $135^\circ$ . You can check to see that your answer is correct:  
 $90 + 90 + 90 + 135 + 135 = 540$ .

## ► Lesson 10

- 1.** true  
**2.** false  
**3.** true  
**4.** false  
**5.** true  
**6.** true  
**7.** false  
**8.** true  
**9.** false  
**10.** false  
**11.** 8  
**12.** 12  
**13.**  $65^\circ$

14.  $115^\circ$
15.  $65^\circ$
16. 14
17. 5
18.  $55^\circ$
19.  $125^\circ$
20.  $125^\circ$
21. 26
22. 13
23.  $25^\circ$
24.  $65^\circ$
25.  $58^\circ$
26.  $64^\circ$
27.  $90^\circ$
28.  $58^\circ$
29.  $116^\circ$
30.  $32^\circ$

### ► Lesson 11

1.  $\frac{10}{5} = \frac{2}{1}$
2.  $\frac{5}{10} = \frac{1}{2}$
3.  $\frac{5}{15} = \frac{1}{3}$
4. yes
5. no
6. no
7.  $x = 4$
8.  $y = 44$
9.  $z = 28$
10.  $a = 8$
11.  $b = 14$
12.  $x = 1$
13. yes
14. no
15. no
16. yes
17. SAS
18. AA, SAS, or SSS
19. AA
20. AA
21. none

22. none
23. SAS
24. SSS

### ► Lesson 12

1. 21
2. 36
3. 33
4. 30
5. 22 in.
6. 18 cm
7. 36 ft.
8. 26 cm
9. 80 cm
10. 36 in.
11. 80 in.
12. 63 ft.
13.  $w = 1$  m;  $l = 5$  m
14.  $w = 5$  cm;  $l = 10$  cm
15.  $s = 7$  in.
16.  $s = 8$  yd.
17.  $s = 20$  cm
18.  $s = 10$  cm
19.  $s = 8$  cm
20.  $s = 23$  cm
21. 36 in.
22. 80 ft.
23. 68 ft.
24. 16 packages

### ► Lesson 13

1.  $24 \text{ cm}^2$
2.  $16 \text{ m}^2$
3.  $196 \text{ in.}^2$
4.  $21 \text{ ft.}^2$
5.  $b = 8$  in.;  $h = 7$  in.
6.  $b = 15$  m;  $h = 5$  m
7.  $h = 6.25$  yd.
8.  $b = 15$  cm



- 9.**  $A = 24 \text{ cm}^2$ ;  $b = 8 \text{ cm}$ ;  $h = 3 \text{ cm}$   
**10.**  $A = 41.5 \text{ mm}^2$ ;  $b = 8.3 \text{ mm}$ ;  $h = 5 \text{ mm}$   
**11.**  $b = 6 \text{ cm}$ ;  $h = 10 \text{ cm}$   
**12.**  $b = 7 \text{ ft.}$ ;  $h = 4 \text{ ft.}$   
**13.**  $A = 21 \text{ cm}^2$   
**14.**  $A = 30 \text{ in.}^2$   
**15.**  $A = 35 \text{ ft.}^2$   
**16.**  $A = 169 \text{ in.}^2$   
**17.**  $h = 14 \text{ cm}$   
**18.**  $h = 8 \text{ ft.}$   
**19.**  $b = 17 \text{ in.}$   
**20.**  $A = 132 \text{ m}^2$   
**21.**  $b_1 = 5 \text{ cm}$   
**22.**  $A = 104 \text{ in.}^2$   
**23.**  $h = 6 \text{ in.}$   
**24.**  $A = 45 \text{ in.}^2$   
**25.**  $b_2 = 9 \text{ m}$

### ► Lesson 14

- 1.** 5 in.  
**2.** 4 in.  
**3.** 9 in.  
**4.** 5 in.  
**5.** 4 in.  
**6.** 9 in.  
**7.** 12 edges  
**8.** 6 faces  
**9.** 8 vertices  
**10.**  $22 \text{ in.}^2$   
**11.**  $78 \text{ ft.}^2$   
**12.**  $158 \text{ m}^2$   
**13.**  $200 \text{ cm}^2$   
**14.**  $174 \text{ in.}^2$   
**15.**  $220 \text{ ft.}^2$   
**16.**  $96 \text{ ft.}^2$   
**17.**  $216 \text{ cm}^2$   
**18.**  $24 \text{ in.}^2$   
**19.**  $294 \text{ m}^2$   
**20.**  $201.84 \text{ cm}^2$   
**21.**  $6e^2 = 486$ ;  $e^2 = 81$ ;  $e = 9 \text{ ft.}$

### ► Lesson 15

- 1.**  $432 \text{ in.}^3$   
**2.**  $32 \text{ m}^3$   
**3.**  $343 \text{ cm}^3$   
**4.**  $240 \text{ ft.}^3$   
**5.**  $108 \text{ m}^3$   
**6.**  $100 \text{ ft.}^3$   
**7.**  $120 \text{ cm}^3$   
**8.**  $150 \text{ m}^3$   
**9.**  $36 \text{ in.}^3$   
**10.**  $1,120 \text{ cm}^3$   
**11.**  $576 \text{ ft.}^3$   
**12.**  $680 \text{ m}^3$   
**13.**  $1,000 \text{ m}^3$   
**14.**  $27 \text{ ft.}^3$   
**15.**  $64 \text{ in.}^3$   
**16.**  $729 \text{ cm}^3$   
**17.**  $192 \text{ cm}^3$   
**18.**  $180 \text{ cm}^3$   
**19.**  $400 \text{ in.}^3$   
**20.**  $60 \text{ ft.}^3$   
**21.**  $270 \text{ cm}^3$   
**22.**  $85 \text{ cm}^3$   
**23.**  $128 \text{ in.}^3$   
**24.**  $416 \text{ ft.}^3$   
**25.**  $1 \text{ cm}^3$


### ► Lesson 16

- 1.** 47.1 ft.  
**2.** 69.08 in.  
**3.** 21.98 m  
**4.** 157 cm  
**5.**  $78.5 \text{ ft.}^2$   
**6.**  $200.96 \text{ m}^2$   
**7.**  $530.66 \text{ cm}^2$   
**8.**  $452.16 \text{ in.}^2$   
**9.**  $81.64 \text{ m}^2$   
**10.**  $747.32 \text{ m}^2$   
**11.**  $828.96 \text{ ft.}^2$

- 12.**  $678.24 \text{ cm}^2$   
**13.**  $1,271.7 \text{ cm}^3$   
**14.**  $339.12 \text{ m}^3$   
**15.**  $1,884 \text{ in.}^3$   
**16.**  $4,559.28 \text{ ft.}^2$   
**17.**  $100.48 \text{ m}^3$   
**18.**  $340.17 \text{ m}^3$   
**19.**  $84.78 \text{ cm}^3$   
**20.**  $256.43 \text{ ft.}^3$   
**21.**  $615.44 \text{ m}^2$   
**22.**  $803.84 \text{ in.}^2$   
**23.**  $3,215.36 \text{ cm}^2$   
**24.**  $1,256 \text{ ft.}^2$   
**25.**  $523.3 \text{ m}^3$   
**26.**  $1,436.03 \text{ m}^3$   
**27.**  $7,234.56 \text{ cm}^3$   
**28.**  $381.51 \text{ ft.}^3$

### ► Lesson 17

1-12.



- 13.** II  
**14.** I  
**15.** IV  
**16.** III  
**17.**  $y$ -axis  
**18.**  $x$ -axis  
**19.** I  
**20.** II  
**21.** III

- 22.** IV  
**23.**  $x$ -axis  
**24.**  $y$ -axis  
**25.**  $A(-4, 2)$   
**26.**  $B(1, -5)$   
**27.**  $C(4, 0)$   
**28.**  $D(-6, -4)$   
**29.**  $E(-2, 6)$   
**30.**  $F(0, -2)$   
**31.**  $G(6, -1)$   
**32.**  $H(3, 4)$   
**33.** 4  
**34.** 10  
**35.** 5  
**36.** 13


### ► Lesson 18

- 1.**  $\frac{1}{3}$   
**2.**  $\frac{2}{-1} = -2$   
**3.**  $\frac{4}{-1} = -4$   
**4.**  $\frac{3}{2}$   
**5.**  $\frac{2}{8} = \frac{1}{4}$   
**6.**  $\frac{2}{-5}$   
**7.** positive  
**8.** negative  
**9.** zero  
**10.** positive  
**11.** undefined (no slope)  
**12.** negative  
**13.**  $\frac{6}{5}$   
**14.** 1  
**15.** 0  
**16.**  $-2$   
**17.** undefined (no slope)  
**18.**  $\frac{3}{4}$   
**19.**  $-\frac{2}{3}$   
**20.** 0

► Lesson 19


1. linear
2. nonlinear
3. nonlinear
4. linear
5. linear
6. nonlinear
7. linear
8. linear
9. yes
10. no
11. no
12. yes
13. yes
14. no
15. yes
16. yes

17.




- (2,7)
- (0,3)
- (-2,-1)

18.




- (2,-2)
- (0,0)
- (-2,2)

19.



- (1,2)
- (0,4)
- (-1,6)

20.



- (1,-1)
- (0,-5)
- (-1,-9)

## ► Lesson 20

1.  $\frac{5}{13} \approx 0.385$
2.  $\frac{12}{13} \approx 0.923$
3.  $\frac{5}{12} \approx 0.417$
4.  $\frac{12}{13} \approx 0.923$
5.  $\frac{5}{13} \approx 0.385$
6.  $\frac{12}{5} = 2.400$
7.  $\frac{7}{25} = 0.280$
8.  $\frac{24}{25} = 0.960$
9.  $\frac{7}{24} \approx 0.292$
10.  $\frac{24}{25} = 0.960$
11.  $\frac{7}{25} = 0.280$
12.  $\frac{24}{7} \approx 3.429$
13. 0.309
14. 0.530
15. 0.891
16. 0.766
17. 0.727
18. 0.306
19.  $29^\circ$
20.  $42^\circ$
21.  $21^\circ$
22.  $38^\circ$
23.  $22^\circ$
24.  $17^\circ$
25.  $45^\circ$
26.  $37^\circ$
27.  $23^\circ$
28.  $30^\circ$
29. 13.2
30. 8.1
31. 7.2
32. 3.4

## ► Posttest

If you miss any of the answers, you can find help for that kind of question in the lesson shown to the right of the answer.

1. c Lesson 3
2. a Lesson 2
3. a Lesson 4
4. b Lesson 5
5. b Lesson 6
6. b Lesson 7
7. b Lesson 8
8. c Lesson 9
9. d Lesson 10
10. a Lesson 11
11. c Lesson 12
12. d Lesson 13
13. c Lesson 14
14. a Lesson 15
15. d Lesson 16
16. c Lesson 17
17. c Lesson 18
18. d Lesson 19
19. c Lesson 20
20. d Lesson 1
21. b Lesson 3
22. c Lesson 2
23. b Lesson 2
24. c Lesson 4
25. d Lesson 4
26. a Lesson 5
27. c Lesson 6
28. c Lesson 6
29. c Lesson 7
30. d Lesson 8
31. c Lesson 8
32. d Lesson 9
33. d Lesson 10
34. b Lesson 10
35. c Lesson 11
36. d Lesson 12

**ANSWER KEY**

**37. c** Lesson 12

**38. c** Lesson 13

**39. b** Lesson 14

**40. b** Lesson 14

**41. b** Lesson 15

**42. d** Lesson 16

**43. c** Lesson 16

**44. b** Lesson 17

**45. b** Lesson 18

**46. b** Lesson 18

**47. a** Lesson 19

**48. a** Lesson 20

**49. a** Lesson 20

**50. d** Lesson 1

# Glossary



**Acute angle:** An angle whose measure is between  $0^\circ$  and  $90^\circ$ .

**Acute triangle:** A triangle with three acute angles.

**Angle:** A figure formed by two rays or two segments with a common endpoint.

**Area:** The amount of surface in a region. Area is expressed in square units.

**Base angles of an isosceles triangle:** The two angles that have the base as part of one side.

**Base of a cone:** Its circular face.

**Base of an isosceles triangle:** In an isosceles triangle with two congruent sides, the third side is called the base of the isosceles triangle.

**Base of a trapezoid:** Either of the parallel sides.

**Circle:** All points in a plane that are the same distance from some point called the center.

**Circumference:** The distance around a circle.

**Collinear points:** Points that lie on a line.

**Complementary angles:** Two angles whose measures total  $90^\circ$ .

**Congruent ( $\cong$ ):** Being the same or equal to.

**Coordinates ( $x,y$ ):** An ordered pair of numbers used to name (locate) a point in a plane.

**Coordinate plane:** A plane that contains a horizontal number line (the  $x$ -axis) and a vertical number line (the  $y$ -axis). Every point in the coordinate plane can be named by a pair of numbers.

**Coplanar points:** Points that lie on a plane.

**Corresponding parts:** Any pair of sides or angles in congruent or similar polygons having the same relative position.

**Cosine (cos):** The ratio of the length of the leg adjacent to an acute angle of a right triangle to the length of the hypotenuse.

**Cube:** A rectangular prism in which all faces are congruent squares.

**Diagonal of a polygon:** A segment that joins two nonconsecutive vertices of the polygon.

**Diameter of a circle:** A segment that joins two points on the circle and passes through the center.

**Edge:** A segment formed by the intersection of two faces of a three-dimensional figure.

**Equiangular triangle:** A triangle with all angles equal.

**Equilateral triangle:** A triangle with all sides equal.

**Extremes:** The first and fourth terms of a proportion.

**Face:** A part of a plane forming a side of a three-dimensional figure.

**Graph of an equation:** The geometric figure that contains all the points whose coordinates make the equation a true statement.

**Hexagon:** A polygon with six sides.

**Hypotenuse:** The side opposite the right angle of a right triangle.

**Intersecting lines:** Lines that meet in one point.

**Isosceles triangle:** A triangle with two congruent sides.

**Legs of a right triangle:** Sides that determine the right angle.

**Line ( $\leftrightarrow$ ):** A line is an undefined term for a set of points that extend indefinitely in two directions.

**Linear equation:** An equation whose graph is a line.

**Means:** The second and third terms of a proportion.

**Measure of an angle:** The number of degrees of an angle. Shown by the notation  $m\angle ABC$ .

**Noncollinear points:** A set of points through which a line cannot be drawn.

**Noncoplanar points:** A set of points through which a plane cannot be drawn.

**Obtuse angle:** An angle whose measure is between  $90^\circ$  and  $180^\circ$ .

**Obtuse triangle:** A triangle with one obtuse angle.

**Octagon:** A polygon with eight sides.

**Opposite rays:** Two rays that have a common endpoint and form a line.

**Parallel lines ( $\parallel$ ):** Coplanar lines that do not intersect.

**Parallelogram ( $\square$ ):** A quadrilateral with two pairs of opposite sides parallel. Rectangles, rhombuses, and squares are special types of parallelograms.

**Pentagon:** A polygon with five sides.

**Perimeter:** The distance around a two-dimensional figure.

**Perpendicular ( $\perp$ ):** Lines, segments, or rays that intersect to form right angles.

**Pi ( $\pi$ ):** The ratio of the circumference of a circle to its diameter. The most commonly used approximations for  $\pi$  are  $\frac{22}{7}$  and 3.14.

**Plane:** An undefined term for a flat surface that extends indefinitely in all directions.

**Point:** An undefined term for a figure that indicates a definite location.

**Polygon:** A simple, closed, two-dimensional figure formed only by line segments that meet at points called vertices.

**Polyhedron:** A three-dimensional figure in which each surface is a polygon.

**Postulate:** A statement that is accepted without proof.

**Proof:** A convincing argument.

**Proportion:** A statement that two ratios are equal.

**Protractor:** An instrument used to measure angles.

**Pythagorean theorem:** For any right triangle, the square of the length of the hypotenuse is equal to the sum of the squares of the lengths of the legs.

**Quadrant:** One of the four regions, labeled I–IV, into which the coordinate axes divide a coordinate plane.

**Quadrilateral:** A polygon with four sides.

**Radius of a circle:** A segment joining the center of the circle to any point on the circle.

**Radius of a sphere:** A segment joining the center of the sphere to any point on the sphere.

**Ratio:** A comparison of two numbers by division. If  $b \neq 0$ , the ratio of  $a$  to  $b$  is denoted by  $\frac{a}{b}$ ,  $a:b$ , or  $a$  to  $b$ .

**Ray ( $\rightarrow$ ):** Part of a line with one endpoint. It extends indefinitely in one direction. A ray is named by its endpoint and any other point on the ray. The endpoint is named first.

**Rectangle:** A parallelogram with four right angles. A square is a special type of rectangle.

**Regular polygon:** A polygon in which all sides are congruent and all angles are congruent.

**Rhombus:** A parallelogram with four congruent sides. A square is a special type of rhombus.

**Right angle:** An angle whose measure is  $90^\circ$ .

**Right triangle:** A triangle that contains one right angle.

**Rise:** The vertical distance from a given point to a second given point.

**Run:** The horizontal distance from a given point to a second given point.

**Scalene triangle:** A triangle with no sides equal.

**Segment:** Part of a line with two endpoints. A segment is named by its endpoints.

**Sides of an angle:** The rays that meet to form an angle.

**Sine (sin):** The ratio of the length of the leg opposite an acute angle of a right triangle to the length of the hypotenuse.

**Skew lines:** Two lines that are not coplanar and do not intersect.

**Slope of a line:** The measure of the steepness of a line. It is also the ratio of rise to run. Also, if a line contains points  $(x_1, y_1)$  and  $(x_2, y_2)$ , then the slope of the line is  $\frac{y_2 - y_1}{x_2 - x_1}$ , provided  $x_1 \neq x_2$ .

**Sphere:** A figure in space whose points are the same distance from a particular point. This point is called the center.

**Square:** A parallelogram with four right angles and four congruent sides. A square can also be defined as a parallelogram that is both a rectangle and a rhombus.

**Square of a number ( $n^2$ ):** A number multiplied by itself.

**Square root ( $\sqrt{\quad}$ ):** The positive number which when squared gives the original number as a product. The square root of 25 is 5 because  $5 \times 5 = 25$ .

**Straight angle:** An angle whose measure is  $180^\circ$ .

**Supplementary angles:** Two angles whose measures total  $180^\circ$ .

**Tangent (tan):** The ratio of the length of the leg opposite an acute angle of a right triangle to the adjacent leg.

**Theorem:** A statement that can be proved.

**Transversal:** A line that intersects two or more coplanar lines at different points.

**Trapezoid:** A quadrilateral with exactly one pair of opposite sides parallel.

**Triangle:** A polygon with three sides.

**Trigonometry:** Mathematical principles based on the properties of similar right triangles.

**Vertex:** The intersection of two sides of a polygon.

**Vertical angles:** Two angles whose sides form two pairs of opposite rays. Vertical angles are congruent.

**Volume:** The amount of space in a solid. Volume is expressed in cubic units.





## A P P E N D I X



# Postulates and Theorems

## ► Postulates

### Chapter 1

- Two points determine exactly one line.
- Three noncollinear points determine exactly one plane.

### Chapter 3

- If two parallel lines are cut by a transversal, then corresponding angles are congruent.

### Chapter 4

- If point  $B$  is in the interior of  $\angle AOC$ , then  $m\angle AOB + m\angle BOC = m\angle AOC$ . (see page 41 for diagram)
- If  $\angle AOC$  is a straight line, then  $m\angle AOB + m\angle BOC = 180^\circ$ . (see page 41 for diagram)

### Chapter 7

- If three sides of one triangle are congruent with three sides of another triangle, then the two triangles are congruent (SSS postulate).

- If two sides and the included angle of one triangle are congruent to the corresponding parts of another triangle, then the triangles are congruent (SAS postulate).
- If two angles and the included side of one triangle are congruent to corresponding parts of another triangle, the triangles are congruent (ASA postulate).

### Chapter 11

- If two angles of one triangle are congruent to two angles of another triangle, then the triangles are similar (AA postulate).
- If the lengths of the corresponding sides of two triangles are proportional, then the triangles are similar (SSS postulate).
- If the lengths of two pairs of corresponding sides of two triangles are proportional and the corresponding included angles are congruent, then the triangles are similar (SAS postulate).

## ► Theorems

### Chapter 3

- If two parallel lines are cut by a transversal, then alternate interior angles are congruent.
- If two parallel lines are cut by a transversal, then same-side interior angles are supplementary.

### Chapter 5

- Vertical angles are congruent.

### Chapter 8

- Pythagorean theorem: In a right triangle, the sum of the squares of the lengths of the legs is equal to the square of the length of the hypotenuse ( $a^2 + b^2 = c^2$ ).
- Converse of the Pythagorean theorem: If the square of the length of the longest side of a

triangle is equal to the sum of the squares of the lengths of the two shorter sides, then the triangle is a right triangle.

- If the square of the length of the longest side is greater than the sum of the squares of the lengths of the other two shorter sides, then the triangle is obtuse ( $c^2 > a^2 + b^2$ ). (see page 73 for diagram)
- If the square of the length of the longest side is less than the sum of the squares of the lengths of the two other sides, then the triangle is acute ( $c^2 < a^2 + b^2$ ). (see page 73 for diagram)

### Chapter 9

- The sum of interior angles of a triangle is  $180^\circ$ .
- If a convex polygon has  $n$  sides, then its angle sum is given by the formula  $S = 180(n - 2)$ .
- The sum of exterior angles of a convex polygon is always  $360^\circ$ .

### Chapter 10

- Opposite sides of a parallelogram are congruent.
- Opposite angles of a parallelogram are congruent.
- Consecutive angles of a parallelogram are supplementary.
- Diagonals of a parallelogram bisect each other.
- Diagonals of a rectangle are congruent.
- The diagonals of a rhombus are perpendicular, and they bisect the angles of the rhombus.

### Chapter 13

- The area ( $A$ ) of a rectangle is the product of its base length ( $b$ ) and its height ( $h$ ):  $A = bh$ .
- The area of a parallelogram ( $A$ ) is the product of its base length ( $b$ ) and its height ( $h$ ):  $A = bh$ .
- The area ( $A$ ) of any triangle is half the product of its base length ( $b$ ) and its height ( $h$ ):  $A = \frac{1}{2}bh$ .
- The area of a trapezoid is half the product of the height and the sum of the base lengths ( $b_1 + b_2$ ):  $A = \frac{1}{2}bh(b_1 + b_2)$ .

**Chapter 14**

- The surface area (S.A.) of a rectangular prism is twice the sum of the length ( $l$ ) times the width ( $w$ ), the width ( $w$ ) times the height ( $h$ ), and the length ( $l$ ) times the height ( $h$ ):  $S.A. = 2(lw + wh + lh)$ .
- The surface area of a cube is six times the edge ( $e$ ) squared:  $S.A. = 6e^2$ .

**Chapter 15**

- To find the volume ( $V$ ) of a rectangular prism, multiply the length ( $l$ ) by the width ( $w$ ) and by the height ( $h$ ):  $V = lwh$ .
- To find the volume ( $V$ ) of any prism, multiply the area of the base ( $B$ ) by the height ( $h$ ):  $V = Bh$ .
- The volume of a cube is determined by cubing the length of the edge:  $V = e^3$ .

**Chapter 16**

- The circumference of any circle is the product of its diameter and  $\pi$ :  $C = \pi d$  or  $C = 2\pi r$ .
- The area ( $A$ ) of a circle is the product of and the square of the radius ( $r$ ):  $A = \pi r^2$ .
- The surface area (S.A.) of a cylinder is determined by finding the sum of the area of the bases and the product of the circumference times the height:  $S.A. = 2\pi r^2 + 2\pi rh$ .
- The volume ( $V$ ) of a cylinder is the product of the area of the base ( $B$ ) and the height ( $h$ ):  $V = Bh$  or  $V = \pi r^2 h$ .
- The surface area (S.A.) formula for a sphere is four times  $\pi$  times the radius squared:  $S.A. = 4\pi r^2$ .
- The volume ( $V$ ) of a sphere is determined by the product of  $\frac{4}{3}\pi$  times the radius cubed:  $V = \frac{4}{3}\pi r^3$ .

**Chapter 17**

- The distance  $d$  between any two points  $A(x_1, y_1)$  and  $B(x_2, y_2)$  is  $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$ .



## A P P E N D I X

# B



# Additional Resources

**M**any resources are available to help you if you need additional practice with geometry. Your local high school is a valuable resource. Most high school math teachers would assist you if you asked them for help with a lesson. Also, a teacher may provide you with practice sets of problems on a lesson that proved difficult for you. If you need a tutor, the teacher may be able to suggest one for you. You could also check the classified ads in your local newspaper or the yellow pages to search for a tutor.

Colleges are also a valuable resource. They often have learning centers or tutor programs available. A college may be another good source for locating a tutor. To find out what is available in your community, call your local college's math department or learning center.

If you would like to continue working geometry problems on your own, your local bookstore or library has books that can help you. You may also be able to borrow a textbook from your local high school. The geometry books listed on the next page provide helpful explanations or practice sets of problems. Check your local bookstore or library for availability.

## APPENDIX B: ADDITIONAL RESOURCES

Arnone, Wendy. *Geometry for Dummies*. (New York: For Dummies) 2001.

Jacobs, Harold R. *Geometry, Third Edition: Seeing, Doing, Understanding*. (New York: W.H. Freeman) 2003.

Long, Lynette. *Painless Geometry*. (Hauppauge: Barron's Educational Series) 2001.

Rich, Barnett. *Schaum's Outline of Geometry, Third Edition*. (New York: McGraw-Hill) 1999.

Slavin, Steve, and Ginny Crisonino. *Geometry: A Self-Teaching Guide*. (New York: Wiley, John & Sons) 2004.